

Groups acting freely on Λ -trees

Olga Kharlampovich*

Alexei Myasnikov[†]

Denis Serbin

September 23, 2011

Abstract

A group is called Λ -free if it has a free Lyndon length function in an ordered abelian group Λ , which is equivalent to having a free isometric action on a Λ -tree. A group has a regular free length function in Λ if and only if it has a free isometric action on a Λ -tree so that all branch points belong to the orbit of the base point. In this paper we prove that every finitely presented Λ -free group G can be embedded into a finitely presented group with a regular free length function in Λ so that the length function on G is preserved by the embedding. Next, we prove that every finitely presented group $\tilde{G}$ with a regular free Lyndon length function in Λ has a regular free Lyndon length function in $\mathbb{R}^n$ ordered lexicographically for an appropriate n and can be obtained from a free group by a series of finitely many HNN-extensions in which associated subgroups are maximal abelian and length isomorphic.

1 Introduction

This is the fourth paper in a series, where we investigate non-Archimedean group actions, length functions and infinite words. In [28] Lyndon introduced real-valued length functions as a tool to carry over Nielsen cancellation theory from free groups to a more general setting (see also [29]). Some results in this direction were obtained in [20, 21, 19, 34, 2]. In [6] Chiswell described a crucial construction which shows that a group with a real-valued length function has an action on an $\mathbb{R}$ -tree, and vice versa. Later, Morgan and Shalen realized that a similar construction holds for an arbitrary group with a Lyndon length function which takes values in an arbitrary ordered abelian group Λ (see [33]). In particular, they introduced Λ -trees as a natural generalization of $\mathbb{R}$ -trees which they studied in relation with Thurston's Geometrization Program. Thus, actions on Λ -trees and Lyndon length functions with values in Λ are two equivalent languages describing the same class of groups. In the case when the action is free (the stabilizer of every point is trivial) we call groups in this class Λ -free. We refer to the book [7] for a detailed discussion on the subject.

One of the major events in combinatorial group theory in 1970's was the development of Bass-Serre theory. We refer to the book [36], where Serre laid down fundamentals of the theory of groups acting freely on simplicial trees. In particular, Bass-Serre theory makes it possible to extract information about the structure of a group from its action on a simplicial tree. Alperin and Bass [1] developed the initial framework of the theory of group actions on Λ -trees and stated the fundamental research goals: find the group theoretic information carried by an action (by isometries) on a Λ -tree; generalize Bass-Serre theory to actions on arbitrary Λ -trees.

A joint effort of several researchers culminated in a description of finitely generated groups acting freely on $\mathbb{R}$ -trees [4, 13], which is now known as Rips' theorem: a finitely generated group acts freely on an $\mathbb{R}$ -tree if and only if it is a free product of free abelian groups and surface groups (with an exception of non-orientable surfaces of genus 1, 2, and 3). The key ingredient of this theory is the

*Supported by NSERC grant

[†]Supported by NSERC grant

so-called “Rips machine”, the idea of which comes from Makanin’s algorithm for solving equations in free groups (see [30]). The Rips machine appears in applications as a general tool that takes a sequence of isometric actions of a group G on some “negatively curved spaces” and produces an isometric action of G on an $\mathbb{R}$ -tree as the Gromov-Hausdorff limit of the sequence of spaces. Free actions on $\mathbb{R}$ -trees cover all Archimedean actions, since every group acting freely on a Λ -tree for an Archimedean ordered abelian group Λ also acts freely on an $\mathbb{R}$ -tree.

In the non-Archimedean case there are only partial results for particular choices of Λ . First of all, in [3] Bass studied finitely generated groups acting freely on $\Lambda_0 \oplus \mathbb{Z}$ -trees with respect to the right lexicographic order on $\Lambda_0 \oplus \mathbb{Z}$. In this case it was shown that the group acting freely on a $\Lambda_0 \oplus \mathbb{Z}$ -tree splits into a graph of groups with Λ_0 -free vertex groups and maximal abelian edge groups. Next, Guirardel (see [18]) obtained the structure of finitely generated groups acting freely on $\mathbb{R}^n$ -trees (with the lexicographic order). In [25] the authors described the class of finitely generated groups acting freely and regularly on $\mathbb{Z}^n$ -trees in terms of HNN-extensions of a very particular type. The importance of regular actions becomes clear from the results of [26], where we proved that a finitely generated group acting freely on a $\mathbb{Z}^n$ -tree is a subgroup of a finitely generated group acting freely and regularly on a $\mathbb{Z}^m$ -tree for $m \geq n$, and the paper [9], where it was shown that a group acting freely on a Λ -tree (for arbitrary Λ) can always be embedded in a length-preserving way into a group acting freely and regularly on a Λ -tree.

In this paper we give a partial solution (for finitely presented groups) of the following main problem of the Alperin-Bass program.

Problem. *Describe finitely presented (finitely generated) Λ -free groups for an arbitrary ordered abelian group Λ .*

Nowadays, the geometric method of group actions seems much more in use than the length functions, even though the Nielsen theory still gives very powerful results (see the discussion in [25]). There are several reasons for this. Firstly, the Lyndon’s abstract axiomatic approach to length functions is less intuitive than group actions. Secondly, the current development of the abstract Nielsen theory is incomplete and insufficient — one of the principal notions of complete actions and regular length functions were not in use until recently. Thirdly, and this is crucial, an abstract analog of the Rips machine for length functions with values in $\mathbb{R}$ (or an arbitrary Λ) was not introduced or developed enough to be used easily in applications. Notice, that in the case of $\Lambda = \mathbb{Z}$ the completeness and regularity come for free, so there was no need for axiomatic formalization and it went mostly unnoticed. The regularity axiom appeared first in [31, 32] as a tool to deal with length functions in $\mathbb{Z}^n$ (with respect to the lexicographic order). On the other hand, an analog of the Rips machine for $\Lambda = \mathbb{Z}$ did exist for a long time — the original Makanin-Razborov process [35] for solving equations in free groups. But it was not recognized as such until [23, 24], where it was used systematically to get splittings of groups.

Introduction of infinite Λ -words in [32] gives another approach to group actions. Every group admitting a faithful representation by Λ -words acts freely on a Λ -tree. The converse is proved in [6]. Formulation of the regularity axiom is another important ingredient. This axiom ensures that the group is complete with respect to the Gromov’s inner product, that is, the Gromov’s product of two elements of the group can be realized as the length of a particular element of the group (not only as an abstract length). In the language of actions this means that all branch points belong to the orbit of the base point. This allows one to use the Nielsen cancelation argument, very similar to the case of a free group. The regularity (completeness) condition is crucial for the existence of Makanin-Razborov type processes over groups with Λ -length functions.

In this paper we prove the following results.

Theorem 1. *Any finitely presented regular Λ -free group G can be represented as a union of a finite series of groups*

$$G_1 < G_2 < \cdots < G_n = G,$$

where

1. G_1 is a free group,
2. G_{i+1} is obtained from G_i by finitely many HNN-extensions in which associated subgroups are maximal abelian and the associated isomorphisms preserve the length induced from G_i .

Theorem 2. *Let G be a finitely presented regular Λ -free group. Then G is a torsion-free total relatively hyperbolic biautomatic group with a quasi-convex hierarchy (see [39] for the definition of a quasi-convex hierarchy).*

Proof. It follows from the description of the structure of G that each G_i is total relatively hyperbolic. (It also follows from [18] and Theorem 3 since G acts freely on $\mathbb{R}^n$ -tree.) From [38] it follows that G is biautomatic. Now by [5] centralizers are regular and quasi-convex in biautomatic groups. So the associated subgroups in G_i are quasiconvex in G_i . If they are not conjugate, then they are centralizers in G_{i+1} and therefore quasiconvex there. If they are conjugate, then the HNN-extension is equivalent to an extension of the centralizer of one of them (denote it by A). The centralizer $C = \langle A, t \mid [a, t] = 1, a \in A \rangle$ is quasiconvex in G_{i+1} and A is a direct summand of C , therefore A is quasiconvex in C and in G_{i+1} . $\square$

Corollary 1. *Let G be a finitely presented regular Λ -free group. Then the following algorithmic problems are decidable in G :*

- the Word and Conjugacy Problems;
- the Diophantine Problem (decidability of arbitrary equations in G).

Indeed, decidability of equations follows from [10]. Results of Dahmani and Groves [11] imply the following two corollaries.

Corollary 2. *Let G be a finitely presented regular Λ -free group. Then:*

- G has a non-trivial abelian splitting and one can find such a splitting effectively,
- G has a non-trivial abelian JSJ-decomposition and one can find such a decomposition effectively.

Corollary 3. *The isomorphism problem is decidable in the class of finitely presented groups that act regularly and freely on some Λ -tree.*

Theorem 3. *Any finitely presented regular Λ -free group G is $\mathbb{R}^n$ -free for an appropriate $n \in \mathbb{N}$, where $\mathbb{R}^n$ is ordered lexicographically.*

Theorem 4. *Any finitely presented Λ -free group can be isometrically embedded into a finitely presented regular Λ -free group.*

Corollary 4. *Any finitely presented Λ -free group G is $\mathbb{R}^n$ -free for an appropriate $n \in \mathbb{N}$ and lexicographically ordered $\mathbb{R}^n$.*

Our main conjecture about Λ -free groups is still open.

Conjecture 1. *Any finitely presented Λ -free group G is $\mathbb{Z}^k$ -free for an appropriate $k \in \mathbb{N}$ and lexicographically ordered $\mathbb{Z}^k$.*

2 Length functions, actions, infinite words

Here we introduce basic definitions and notations which are to be used throughout the whole paper.

2.1 Lyndon length functions and free actions

Let G be a group and Λ an ordered abelian group. Then a function $l : G \rightarrow \Lambda$ is called a (*Lyndon*) *length function* on G if the following conditions hold:

- (L1) $\forall g \in G : l(g) \geq 0$ and $l(1) = 0$;
- (L2) $\forall g \in G : l(g) = l(g^{-1})$;
- (L3) $\forall g, f, h \in G : c(g, f) > c(g, h) \rightarrow c(g, h) = c(f, h)$,
where $c(g, f) = \frac{1}{2}(l(g) + l(f) - l(g^{-1}f))$.

Sometimes we refer to length functions with values in Λ as to Λ -length functions. Observe, that the word length function on a finitely generated group (with respect to a fixed finite generating set) usually is not a Lyndon length function. In this paper we consider length functions only of Lyndon type.

Notice that $c(g, f)$ may not be defined in Λ (if $l(g) + l(f) - l(g^{-1}f)$ is not divisible by 2), so in the axiom (L3) we assume that Λ is canonically embedded into a divisible ordered abelian group $\Lambda_{\mathbb{Q}} = \Lambda \otimes_{\mathbb{Z}} \mathbb{Q}$ (see [32] for details).

It is not difficult to derive the following two properties of length functions from the axioms (L1)–(L3):

- $\forall g, f \in G : l(gf) \leq l(g) + l(f)$;
- $\forall g, f \in G : 0 \leq c(g, f) \leq \min\{l(g), l(f)\}$.

A length function $l : G \rightarrow \Lambda$ is called *free* if it satisfies the following two axioms.

- (L4) $\forall g, f \in G : c(g, f) \in \Lambda$.
- (L5) $\forall g \in G : g \neq 1 \rightarrow l(g^2) > l(g)$.

As we have mentioned in the introduction group actions on Λ -trees provide a geometric counterpart to Λ -length functions. To explain we need the following definitions, which we also use later in the text.

Let X be a non-empty set and Λ an ordered abelian group. A Λ -metric on X is a mapping $p : X \times X \rightarrow \Lambda$ such that for all $x, y, z \in X$:

- (M1) $p(x, y) \geq 0$ and $p(x, y) = 0$ if and only if $x = y$,
- (M2) $p(x, y) = p(y, x)$,
- (M3) $p(x, y) \leq p(x, z) + p(y, z)$.

If the axioms (M1)–(M3) are satisfied then the pair (X, p) is an Λ -metric space. If (X, p) and (X', p') are Λ -metric spaces, an *isometry* from (X, p) to (X', p') is a mapping $f : X \rightarrow X'$ such that $p(x, y) = p'(f(x), f(y))$ for all $x, y \in X$.

For elements $a, b \in \Lambda$ the *closed segment* $[a, b]$ is defined by

$$[a, b] = \{c \in \Lambda \mid a \leq c \leq b\}.$$

More generally, a *segment* in an Λ -metric space is the image of an isometry $\alpha : [a, b] \rightarrow X$ for some $a, b \in \Lambda$. The endpoints of the segment are $\alpha(a), \alpha(b)$. A segment with endpoints $x, y \in X$ is denoted by $[x, y]$. An Λ -metric space (X, p) is *geodesic* if for all $x, y \in X$, there is a segment in X with endpoints x, y .

An Λ -tree is an Λ -metric space (X, p) such that:

- (T1) (X, p) is geodesic,
- (T2) if two segments of (X, p) intersect in a single point, which is an endpoint of both, then their union is a segment,
- (T3) the intersection of two segments with a common endpoint is also a segment.

We say that a group G acts on a Λ -tree X if it acts on X by isometries, i.e. there exists an embedding of G into $\text{Isom}(X)$. An action of G on X is termed *free* if for every $1 \neq g \in G$ neither g , nor g^2 has a fixed point in X .

If G acts on a Λ -tree (X, p) and $x \in X$ then one can associate to this action a function $l_x : G \rightarrow \Lambda$ so that $l_x(g) = p(x, gx)$ for every $g \in G$. It is not hard to check that l_x is a Lyndon length function. Moreover, if the action of G on X is free then l_x is a free length function. The converse is also true, that is, every action of a group G on a Λ -tree arises from some Lyndon Λ -length function.

Theorem 5. [6, 33] *Let G be a group and $l : G \rightarrow \Lambda$ a Lyndon length function satisfying (L4). Then there are a Λ -tree (X, p) , an action of G on X and a point $x \in X$ such that $l = l_x$.*

For elements $g_1, \dots, g_n \in G$ we write

$$g = g_1 \circ \dots \circ g_n$$

if $g = g_1 \dots g_n$ and $l(g) = l(g_1) + \dots + l(g_n)$. Also, for $\alpha \in \Lambda$ we write $g = g_1 \circ_\alpha g_2$ if $g = g_1 g_2$ and $c(g_1^{-1}, g_2) < \alpha$.

A length function $l : G \rightarrow \Lambda$ is called *regular* if it satisfies the *regularity* axiom:

- (L6) $\forall g, f \in G, \exists u, g_1, f_1 \in G$:

$$g = u \circ g_1 \ \& \ f = u \circ f_1 \ \& \ l(u) = c(g, f).$$

Many examples of groups with regular free length functions are given in [25].

2.2 Regular actions

In this subsection we give a geometric characterization of group actions that come from regular length functions. The four lemmas below were proved in [25], but for convenience of the reader we give the proofs here too.

Definition 1. *Let G act on a Λ -tree Γ . The action is regular with respect to $x \in \Gamma$ if for any $g, h \in G$ there exists $f \in G$ such that $[x, fx] = [x, gx] \cap [x, hx]$.*

Lemma 1. [25] *Let G act on a Λ -tree (Γ, d) . Then the action of G is regular with respect to $x \in \Gamma$ if and only if the length function $l_x : G \rightarrow \Lambda$ based at x is regular.*

Proof. By definition, the length function l_x is regular if for every $g, h \in G$ there exists $f \in G$ such that $g = fg_1$, $h = fh_1$, where $l_x(f) = c(g, h)$ and $l_x(g) = l_x(f) + l_x(g_1)$, $l_x(h) = l_x(f) + l_x(h_1)$.

Suppose the action of G is regular with respect to x . Then for $g, h \in G$ there exists $f \in G$ such that $[x, fx] = [x, gx] \cap [x, hx]$. We have $[x, gx] = [x, fx] \cup [fx, gx]$, $[x, hx] = [x, fx] \cup [fx, hx]$ and $[fx, gx] = [x, (f^{-1}g)x] = l_x(f^{-1}g)$, $[fx, hx] = [x, (f^{-1}h)x] = l_x(f^{-1}h)$. Taking $g_1 = f^{-1}g$, $h_1 = f^{-1}h$ we have $l_x(g) = l_x(f) + l_x(g_1)$, $l_x(h) = l_x(f) + l_x(h_1)$. Finally, since $c(g, h) = \frac{1}{2}(l_x(g) + l_x(h) - l_x(g^{-1}h))$ and $l_x(g^{-1}h) = d(x, (g^{-1}h)x) = d(gx, hx) = d(fx, gx) + d(fx, hx)$ we get $l_x(f) = c(g, h)$.

Suppose that l_x is regular. Then from $g = f \circ g_1$, $h = f \circ h_1$ it follows that $[x, gx] = [x, fx] \cup [fx, gx]$, $[x, hx] = [x, fx] \cup [fx, hx]$. $l_x(f) = c(g, h) = \frac{1}{2}(l_x(g) + l_x(h) - l_x(g^{-1}h))$, so $2d(x, fx) = d(x, gx) + d(x, hx) - d(x, (g^{-1}h)x) = d(x, gx) + d(x, hx) - d(gx, hx)$. In other words,

$$d(gx, hx) = d(x, gx) + d(x, hx) - 2d(x, fx) = (d(x, gx) - d(x, fx)) + (d(x, hx) - d(x, fx))$$

$$= d(fx, gx) + d(fx, hx)$$

which is equivalent to $[x, fx] = [x, gx] \cap [x, hx]$. $\square$

Lemma 2. *Let G act minimally on a Λ -tree Γ . If the action of G is regular with respect to $x \in \Gamma$ then all branch points of Γ are G -equivalent.*

Proof. From minimality of the action it follows that Γ is spanned by the set of points $Gx = \{gx \mid g \in G\}$.

Now let y be a branch point in Γ . It follows that there exist (not unique in general) $g, h \in G$ such that $[x, y] = [x, gx] \cap [x, hx]$. From regularity of the action it follows that there exists $f \in G$ such that $y = fx$. Hence, every branch point is G -equivalent to x and the statement of the lemma follows. $\square$

Lemma 3. *Let G act on a Λ -tree Γ . If the action of G is regular with respect to $x \in \Gamma$ then it is regular with respect to any $y \in Gx$.*

Proof. We have to show that for every $g, h \in G$ there exists $f \in G$ such that $[y, fy] = [y, gy] \cap [y, hy]$. Since $y = tx$ for some $t \in G$ then we have to prove that $[tx, (ft)x] = [tx, (gt)x] \cap [tx, (ht)x]$. The latter equality is equivalent to $[x, (t^{-1}ft)x] = [x, (t^{-1}gt)x] \cap [x, (t^{-1}ht)x]$ which follows from regularity of the action with respect to x . $\square$

Lemma 4. *Let G act freely on a Λ -tree Γ so that all branch points of Γ are G -equivalent. Then the action of G is regular with respect to any branch point in Γ .*

Proof. Let x be a branch point in Γ and $g, h \in G$. If $g = h$ then $[x, gx] \cap [x, hx] = [x, gx]$ and g is the required element. Suppose $g \neq h$. Since the action is free then $gx \neq hx$ and we consider the tripod formed by x, gx, hx . Hence, $y = Y(x, gx, hx)$ is a branch point in Γ and by the assumption there exists $f \in G$ such that $y = fx$. $\square$

2.3 Infinite words and length functions

In this subsection at first we recall some notions from the theory of ordered abelian groups (for all the details refer to the books [16] and [27]) and then following [32] describe the construction of infinite words.

Let Λ be a discretely ordered abelian group with the minimal positive element 1. It is going to be clear from the context if we are using 1 as an element of Λ , or as an integer. Let $X = \{x_i \mid i \in I\}$ be a set. Put $X^{-1} = \{x_i^{-1} \mid i \in I\}$ and $X^\pm = X \cup X^{-1}$. A Λ -word is a function of the type

$$w : [1, \alpha_w] \rightarrow X^\pm,$$

where $\alpha_w \in \Lambda$, $\alpha_w \geq 0$. The element α_w is called the *length* $|w|$ of w .

By $W(\Lambda, X)$ we denote the set of all Λ -words. Observe, that $W(\Lambda, X)$ contains an empty Λ -word which we denote by ε .

Concatenation uv of two Λ -words $u, v \in W(\Lambda, X)$ is a Λ -word of length $|u| + |v|$ and such that:

$$(uv)(a) = \begin{cases} u(a) & \text{if } 1 \leq a \leq |u| \\ v(a - |u|) & \text{if } |u| < a \leq |u| + |v| \end{cases}$$

A Λ -word w is *reduced* if $w(\beta + 1) \neq w(\beta)^{-1}$ for each $1 \leq \beta < |w|$. We denote by $R(\Lambda, X)$ the set of all reduced Λ -words. Clearly, $\varepsilon \in R(\Lambda, X)$. If the concatenation uv of two reduced Λ -words u and v is also reduced then we write $uv = u \circ v$.

For $u \in W(\Lambda, X)$ and $\beta \in [1, \alpha_u]$ by u_β we denote the restriction of u on $[1, \beta]$. If $u \in R(\Lambda, X)$ and $\beta \in [1, \alpha_u]$ then

$$u = u_\beta \circ \tilde{u}_\beta,$$

for some uniquely defined $\tilde{u}_\beta$.

An element $\text{com}(u, v) \in R(\Lambda, X)$ is called the (*longest*) *common initial segment* of Λ -words u and v if

$$u = \text{com}(u, v) \circ \tilde{u}, \quad v = \text{com}(u, v) \circ \tilde{v}$$

for some (uniquely defined) Λ -words $\tilde{u}, \tilde{v}$ such that $\tilde{u}(1) \neq \tilde{v}(1)$.

Now, we can define the product of two Λ -words. Let $u, v \in R(\Lambda, X)$. If $\text{com}(u^{-1}, v)$ is defined then

$$u^{-1} = \text{com}(u^{-1}, v) \circ \tilde{u}, \quad v = \text{com}(u^{-1}, v) \circ \tilde{v},$$

for some uniquely defined $\tilde{u}$ and $\tilde{v}$. In this event put

$$u * v = \tilde{u}^{-1} \circ \tilde{v}.$$

The product $*$ is a partial binary operation on $R(\Lambda, X)$.

An element $v \in R(\Lambda, X)$ is termed *cyclically reduced* if $v(1)^{-1} \neq v(|v|)$. We say that an element $v \in R(\Lambda, X)$ admits a *cyclic decomposition* if $v = c^{-1} \circ u \circ c$, where $c, u \in R(\Lambda, X)$ and u is cyclically reduced. Observe that a cyclic decomposition is unique (whenever it exists). We denote by $CR(\Lambda, X)$ the set of all cyclically reduced words in $R(\Lambda, X)$ and by $CDR(\Lambda, X)$ the set of all words from $R(\Lambda, X)$ which admit a cyclic decomposition.

Let I_Λ index the set of all convex subgroups of Λ . I_Λ is linearly ordered (see, for example, [7]): $i < j$ if and only if $\Lambda_i < \Lambda_j$, and

$$\Lambda = \bigcup_{i \in I_\Lambda} \Lambda_i.$$

We say that $g \in G$ has the height $i \in I_\Lambda$ and denote $ht(g) = i$ if $|g| \in \Lambda_i$ and $|g| \notin \Lambda_j$ for any $j < i$.

Below we refer to Λ -words as *infinite words* usually omitting Λ whenever it does not produce any ambiguity.

The following result establishes the connection between infinite words and length functions.

Theorem 6. [32] *Let Λ be a discretely ordered abelian group and X be a set. Then any subgroup G of $CDR(\Lambda, X)$ has a free Lyndon length function with values in Λ – the restriction $L|_G$ on G of the standard length function L on $CDR(\Lambda, X)$.*

The converse of the theorem above is also true.

Theorem 7. [8] *Let G have a free Lyndon length function $L : G \rightarrow \Lambda$, where Λ is a discretely ordered abelian group. Then there exists a set X and a length preserving embedding $\phi : G \rightarrow CDR(\Lambda, X)$, that is, $|\phi(g)| = L(g)$ for any $g \in G$.*

Corollary 5. [8] *Let G have a free Lyndon length function $L : G \rightarrow \Lambda$, where Λ is an arbitrary ordered abelian group. Then there exists an embedding $\phi : G \rightarrow CDR(\Lambda', X)$, where $\Lambda' = \mathbb{Z} \oplus \Lambda$ is discretely ordered with respect to the right lexicographic order and X is some set, such that, $|\phi(g)| = (0, L(g))$ for any $g \in G$.*

Theorem 6, Theorem 7, and Corollary 5 show that a group has a free Lyndon length function if and only if it embeds into a set of infinite words and this embedding preserves the length. Moreover, it is not hard to show that this embedding also preserves regularity of the length function.

Theorem 8. [22] *Let G have a free regular Lyndon length function $L : G \rightarrow \Lambda$, where Λ is an arbitrary ordered abelian group. Then there exists an embedding $\phi : G \rightarrow R(\Lambda', X)$, where Λ' is a discretely ordered abelian group and X is some set, such that, the Lyndon length function on $\phi(G)$ induced from $R(\Lambda', X)$ is regular.*

3 Generalized equations

From now on we assume that $G = \langle X \mid R \rangle$ is a finitely presented group which acts freely and regularly on a Λ -tree, where Λ is a discretely ordered abelian group, or, equivalently, G can be represented by Λ -words over some alphabet Z and the length function on G induced from $CDR(\Lambda, Z)$ is regular. Let us fix the embedding $\xi : G \hookrightarrow CDR(\Lambda, Z)$ for the rest of this section.

3.1 The notion of a generalized equation

Definition 2. A combinatorial generalized equation Ω (which is convenient to visualize as shown on the picture below)




Figure 1: A typical generalized equation.

consists of the following objects.

1. A finite set of bases $\mathcal{M} = BS(\Omega)$. The set of bases $\mathcal{M}$ consists of $2n$ elements $\mathcal{M} = \{\mu_1, \dots, \mu_{2n}\}$. The set $\mathcal{M}$ comes equipped with two functions: a function $\varepsilon : \mathcal{M} \rightarrow \{1, -1\}$ and an involution $\Delta : \mathcal{M} \rightarrow \mathcal{M}$ (that is, Δ is a bijection such that Δ^2 is an identity on $\mathcal{M}$). Bases μ and $\bar{\mu} = \Delta(\mu)$ are called dual bases. We denote bases by letters μ, λ , etc.
2. A set of boundaries $BD = BD(\Omega) = \{1, 2, \dots, p+1\}$, that is, integer points of the interval $I = [1, p+1]$. We use letters i, j , etc. for boundaries.
3. Two functions $\alpha : BS \rightarrow BD$ and $\beta : BS \rightarrow BD$. We call $\alpha(\mu)$ and $\beta(\mu)$ the initial and terminal boundaries of the base μ (or endpoints of μ). These functions satisfy the following conditions: for every base $\mu \in BS$: $\alpha(\mu) < \beta(\mu)$ if $\varepsilon(\mu) = 1$ and $\alpha(\mu) > \beta(\mu)$ if $\varepsilon(\mu) = -1$.
4. A set of boundary connections (p, λ, q) , where p is a boundary on λ (that is a number between $\alpha(\lambda)$ and $\beta(\lambda)$) and q on $\bar{\lambda}$. In this case we say that p and q are λ -tied. If (p, λ, q) is a boundary connection then $(q, \bar{\lambda}, p)$ is also a boundary connection. (The meaning of boundary connections will be explained in the transformation (ET5)).

With a combinatorial generalized equation Ω one can canonically associate a system of equations in variables $h = (h_1, \dots, h_p)$ (variables h_i are also called *items*). This system is called a *generalized equation*, and (slightly abusing the terminology) we denote it by the same symbol Ω , or $\Omega(h)$ specifying the variables it depends on. The generalized equation Ω consists of the following two types of equations.

1. Each pair of dual bases $(\lambda, \bar{\lambda})$ provides an equation

$$[h_{\alpha(\lambda)} h_{\alpha(\lambda)+1} \dots h_{\beta(\lambda)-1}]^{\varepsilon(\lambda)} = [h_{\alpha(\bar{\lambda})} h_{\alpha(\bar{\lambda})+1} \dots h_{\beta(\bar{\lambda})-1}]^{\varepsilon(\bar{\lambda})}.$$

These equations are called *basic equations*.

2. Every boundary connection (p, λ, q) gives rise to a *boundary equation*

$$[h_{\alpha(\lambda)}h_{\alpha(\lambda)+1} \cdots h_{p-1}] = [h_{\alpha(\bar{\lambda})}h_{\alpha(\bar{\lambda})+1} \cdots h_{q-1}],$$

if $\varepsilon(\lambda) = \varepsilon(\bar{\lambda})$ and

$$[h_{\alpha(\lambda)}h_{\alpha(\lambda)+1} \cdots h_{p-1}] = [h_q h_{q+1} \cdots h_{\alpha(\bar{\lambda})-1}]^{-1},$$

if $\varepsilon(\lambda) = -\varepsilon(\bar{\lambda})$.

Remark 1. We assume that every generalized equation comes from a combinatorial one.

Given a generalized equation $\Omega(h)$ one can define the *group of $\Omega(h)$*

$$G_\Omega = \langle h \mid \Omega(h) \rangle.$$

Definition 3. Let $\Omega(h) = \{L_1(h) = R_1(h), \dots, L_s(h) = R_s(h)\}$ be a generalized equation in variables $h = (h_1, \dots, h_\rho)$. A set $U = (u_1, \dots, u_\rho) \subseteq R(\Lambda, Z)$ of nonempty Λ -words is called a *solution of Ω* if:

1. all words $L_i(U), R_i(U)$ are reduced,
2. $L_i(U) = R_i(U)$, $i \in [1, s]$.

Observe that a solution U of $\Omega(h)$ defines a homomorphism $\xi_U : G_\Omega \rightarrow R(\Lambda, Z)$ induced by the mapping $h_i \rightarrow u_i$, $i \in [1, \rho]$ since after this substitution all the equations of $\Omega(h)$ turn into identities in $R(\Lambda, Z)$.

If we specify a particular solution U of a generalized equation Ω then we use a pair (Ω, U) .

Definition 4. A *cancellation table* $C(U)$ of a solution $U = (u_1, \dots, u_\rho)$ is defined as follows

$$C(U) = \{h_i^\epsilon h_j^\sigma \mid \text{there is cancelation in the product } u_i^\epsilon * u_j^\sigma, \text{ where } \epsilon, \sigma = \pm 1\}.$$

Definition 5. A solution U^+ of a generalized equation Ω is called *consistent with a solution U* if $C(U^+) \subseteq C(U)$.

3.2 From a finitely presented group to a generalized equation


Recall that $G = \langle X \mid R \rangle$ is finitely presented and let $X = \{x_1, \dots, x_n\}$ and $R = \{r_1(X), \dots, r_m(X)\}$. Adding, if necessary, auxiliary generators, we can assume that every relator involves at most three generators.

Since ξ is a homomorphism it follows that after the substitution $x_i \rightarrow \xi(x_i)$, $i \in [1, n]$ all products $r_i(\xi(X))$, $i \in [1, m]$ cancel out. Hence, we have finitely many *cancellation diagrams* over $CDR(\Lambda, Z)$, which give rise to a generalized equation Ω corresponding to the embedding $\xi : G \hookrightarrow CDR(\Lambda, Z)$.

The precise definition and all the details concerning cancellation diagrams over $CDR(\Lambda, Z)$ can be found in [22]. Briefly, a cancellation diagram for $r_i(\xi(X))$ can be viewed as a finite directed tree T_i in which every positive edge e has a label λ_e so that every occurrence x^δ , $\delta \in \{-1, 1\}$ of $x \in X$ in r_i corresponds to a reduced path $e_1^{\epsilon_1} \cdots e_k^{\epsilon_k}$, where $\epsilon_i \in \{-1, 1\}$, in T_i and $\xi(x^\delta) = \lambda_{e_1}^{\epsilon_1} \circ \cdots \circ \lambda_{e_k}^{\epsilon_k}$. In other words, each λ_e is a piece of some generator of G viewed as a Λ -word. Moreover, we assume that $|\lambda_e|$ is known (since we know the homomorphism ξ).

Now we would like to construct a generalized equation Ω_i corresponding to T_i . Denote by $X(T_i)$ all generators of G which appear in r_i . Next, consider a segment J in Λ of length

$$\sum_{x \in X(T_i)} |\xi(x)|$$

Figure 2: From the cancellation diagram for the relation $[x, y] = 1$ to the generalized equation.

which is naturally divided by the lengths of $\xi(x)$, $x \in X(T_i)$ into subsegments with respect to any given order on $X(T_i)$. Since every $\xi(x)$, $x \in X(T_i)$ splits into at least one reduced product $\lambda_{e_1}^{\epsilon_1} \circ \dots \circ \lambda_{e_k}^{\epsilon_k}$, every such splitting gives a subdivision of the corresponding subsegment of J . Hence, we subdivide J using all product representations of all $\xi(x)$, $x \in X(T_i)$. As a result we obtain a subdivision of J into ρ_i items whose endpoints become boundaries of Ω_i . Observe that each λ_e appears exactly twice in the products representing some $\xi(x)$, $x \in X(T_i)$ and each such entry covers several adjacent items of J . This pair of entries defines a pair of dual bases $(\lambda_e, \bar{\lambda}_e)$. Hence,

$$\mathcal{M}_i = BS(\Omega_i) = \{\lambda_e, \bar{\lambda}_e \mid e \in E(T_i)\}.$$

$\epsilon(\lambda_e)$ depends on the sign of λ_e in the corresponding product representing a variable from $X(T_i)$ (similarly for $\bar{\lambda}_e$).

In the same way one can construct T_i and the corresponding Ω_i for each r_i , $i \in [1, m]$. Combining all combinatorial generalized equations Ω_i , $i \in [1, m]$ we obtain the equation Ω with items $h_1, \dots, h_\rho$ and bases $\mathcal{M} = \cup_i \mathcal{M}_i$. By definition

$$G_\Omega = \langle h_1, \dots, h_\rho \mid \Omega(h_1, \dots, h_\rho) \rangle.$$

At the same time, since each item can be obtained in the form

$$(\lambda_{i_1}^{\epsilon_1} \circ \dots \circ \lambda_{i_k}^{\epsilon_k}) * (\lambda_{j_1}^{\delta_1} \circ \dots \circ \lambda_{j_l}^{\delta_l})^{-1},$$

it follows that G_Ω can be generated by $\mathcal{M}$ with the relators obtained by rewriting $\Omega(h_1, \dots, h_\rho)$ in terms of $\mathcal{M}$.

It is possible to transform the presentation $\langle h_1, \dots, h_\rho \mid \Omega \rangle$ into $\langle X \mid R \rangle$ using Tietze transformations as follows. From the cancellation diagrams constructed for each relator in R it follows that

$x_i = w_i(h_1, \dots, h_\rho) = w_i(\bar{h})$, $i \in [1, n]$. Hence

$$\langle h_1, \dots, h_\rho \mid \Omega \rangle \simeq \langle h_1, \dots, h_\rho, X \mid \Omega \cup \{x_i = w_i(\bar{h}), i \in [1, n]\} \rangle.$$

Next, from the cancellation diagrams it follows that R is a set of consequences of $\Omega \cup \{x_i = w_i(\bar{h}), i \in [1, n]\}$, hence,

$$\langle h_1, \dots, h_\rho \mid \Omega \rangle \simeq \langle h_1, \dots, h_\rho, X \mid \Omega \cup \{x_i = w_i(\bar{h}), i \in [1, n]\} \cup R \rangle.$$

Finally, since the length function on G is regular, for each h_i there exists a word $u_i(X)$ such that $h_i = u_i(\xi(X))$ and all the equations in $\Omega \cup \{x_i = w_i(\bar{h}), i \in [1, n]\}$ follow from R after we substitute h_i by $u_i(X)$ for each i . It follows that

$$\begin{aligned} & \langle h_1, \dots, h_\rho, X \mid \Omega \cup \{x_i = w_i(\bar{h}), i \in [1, n]\} \cup R \rangle \\ & \simeq \langle h_1, \dots, h_\rho, X \mid \Omega \cup \{x_i = w_i(\bar{h}), i \in [1, n]\} \cup R \cup \{h_j = u_j(X), j \in [1, \rho]\} \rangle \simeq \langle X \mid R \rangle. \end{aligned}$$

It follows that $G \simeq G_\Omega$.

Let B be the standard CW 2-complex for G corresponding to the presentation $\langle \mathcal{M} \mid \Omega(\mathcal{M}) \rangle$. Let $C(\Omega)$ be a CW 2-complex obtained from G_Ω by creating for each relator a polygon with boundary labeled by this relator and gluing these polygons together along edges labeled by the same letter. $C(\Omega)$ may have several vertices. If we add arcs joining all the vertices in $C(\Omega)$ we obtain a CW complex $D(\Omega)$ which has the same fundamental group as B (namely, G).

Lemma 5. *G is a free product of $\pi_1(C(\Omega))$ and, possibly, a free group.*

Proof. Follows from the construction. □

Remark 2. *Let $\tilde{G}$ be a finitely presented group with a free length function in Λ (not necessary regular). It can be embedded isometrically in the group $\hat{G}$ with a free regular length function in Λ by [9]. That group can be embedded in $R(\Lambda', X)$. When we make a generalized equation Ω for $\tilde{G}$, we have to add finite number of elements from $\hat{G}$. Let G be a subgroup generated in $\hat{G}$ by $\tilde{G}$ and these elements. Then G is the quotient of G_Ω containing $\tilde{G}$ as a subgroup.*

4 Elimination process

In this section at first we introduce transformations of combinatorial generalized equations, then we show how they can be applied to “simplify” the given equation Ω and to extract information about the structure of the corresponding group G_Ω .

4.1 Elementary transformations

In this subsection we describe *elementary transformations* of generalized equations. Let (Ω, U) be a generalized equation together with a solution U . An elementary transformation (ET) associates to a generalized equation (Ω, U) a generalized equation (Ω_1, U_1) and an epimorphism $\pi : G_\Omega \rightarrow G_{\Omega_1}$ such that for the solution U_1 the following diagram commutes

$$\begin{array}{ccc} G_\Omega & \xrightarrow{\pi} & G_{\Omega_1} \\ \xi_U \downarrow & \searrow \xi_{U_1} & \\ R(\Lambda, Z) & & \end{array}$$

One can view (ET) as a mapping $ET : (\Omega, U) \rightarrow (\Omega_1, U_1)$.

- (ET1) (*Cutting a base (see Fig. 3)*). Let λ be a base in Ω and p an internal boundary of λ (that is, $p \neq \alpha(\lambda), \beta(\lambda)$) with a boundary connection (p, λ, q) . Then we cut the base λ at p into two new bases λ_1 and λ_2 , and cut $\bar{\lambda}$ at q into the bases $\bar{\lambda}_1$ and $\bar{\lambda}_2$.




Figure 3: Elementary transformation (ET1).

- (ET2) (*Transferring a base (see Fig. 4)*). If a base λ of Ω contains a base μ (that is, $\alpha(\lambda) \leq \alpha(\mu) < \beta(\mu) \leq \beta(\lambda)$) and all boundaries on μ are λ -tied by boundary some connections then we transfer μ from its location on the base λ to the corresponding location on the base $\bar{\lambda}$.




Figure 4: Elementary transformation (ET2).

- (ET3) (*Removal of a pair of matched bases (see Fig. 5)*). If the bases λ and $\bar{\lambda}$ are *matched* (that is, $\alpha(\lambda) = \alpha(\bar{\lambda}), \beta(\lambda) = \beta(\bar{\lambda})$) then we remove $\lambda, \bar{\lambda}$ from Ω .




Figure 5: Elementary transformation (ET3).

Remark 3. Observe, that Ω and Ω_1 , where $\Omega_1 = ET_i(\Omega)$ for $i \in \{1, 2, 3\}$ have the same set of variables h and the bijection $h_i \rightarrow h_i$, $i \in [1, \rho]$ induces an isomorphism $G_\Omega \rightarrow G_{\Omega_1}$. Moreover, U is a solution of Ω if and only if U is a solution of Ω_1 .

(ET4) (*Removal of a lone base (see Fig. 6)*). Suppose, a base λ in Ω does not *intersect* any other base, that is, the items $h_{\alpha(\lambda)}, \dots, h_{\beta(\lambda)-1}$ are contained only inside of the base λ .




Figure 6: Elementary transformation (ET4).

Suppose also that all boundaries in λ are λ -tied, that is, for every i ($\alpha(\lambda) < i \leq \beta(\lambda) - 1$) there exists a boundary $b(i)$ such that $(i, \lambda, b(i))$ is a boundary connection in Ω . Then we remove the pair of bases λ and $\bar{\lambda}$ together with all the boundaries $\alpha(\lambda) + 1, \dots, \beta(\lambda) - 1$ (and rename the rest $\beta(\lambda) - \alpha(\lambda) - 1$ of the boundaries correspondingly).

We define the isomorphism $\pi : G_\Omega \rightarrow G_{\Omega_1}$ as follows:

$$\pi(h_j) = h_j \text{ if } j < \alpha(\lambda) \text{ or } j \geq \beta(\lambda)$$

$$\pi(h_i) = \begin{cases} h_{b(i)} \cdots h_{b(i)-1}, & \text{if } \varepsilon(\lambda) = \varepsilon(\bar{\lambda}), \\ h_{b(i)} \cdots h_{b(i)-1}, & \text{if } \varepsilon(\lambda) = -\varepsilon(\bar{\lambda}) \end{cases}$$

for $\alpha + 1 \leq i \leq \beta(\lambda) - 1$.




Figure 7: Elementary transformation (ET5).

(ET5) (*Introduction of a boundary* (see Fig. 7)). Suppose a point p in a base λ is not λ -tied. The transformation (ET5) λ -ties it. To this end, denote by u_λ the element of $CDR(\Lambda, Z)$ corresponding to λ and let u'_λ be the beginning of this word ending at p . Then we perform one of the following two transformations according to where the end of u'_λ on $\bar{\lambda}$ is situated:

- (a) If the end of u'_λ on $\bar{\lambda}$ is situated on the boundary q then we introduce the boundary connection (p, λ, q) . In this case the corresponding isomorphism $\pi : G_\Omega \rightarrow G_{\Omega_1}$ is induced by the bijection $h_i \rightarrow h_i$, $i \in [1, \rho]$. (If we began with the group $\tilde{G}$ with non-regular length function as in Remark 2 this is the only place where $\pi : G_\Omega \rightarrow G_{\Omega_1}$ may be a proper epimorphism, but its restriction on $\tilde{G}$ is still an isomorphism.)
- (b) If the end of u'_λ on $\bar{\lambda}$ is situated between q and $q+1$ then we introduce a new boundary q' between q and $q+1$ (and rename all the boundaries), and also introduce a new boundary connection (p, λ, q') . In this case the corresponding isomorphism $\pi : G_\Omega \rightarrow G_{\Omega_1}$ is induced by the map $\pi(h) = h$, if $h \neq h_q$, and $\pi(h_q) = h_{q'}h_{q'+1}$.

4.2 Derived transformations and auxiliary transformations

In this section we define complexity of a generalized equation and describe several useful “derived” transformations of generalized equations. Some of them can be realized as finite sequences of elementary transformations, others result in equivalent generalized equations but cannot be realized by finite sequences of elementary moves.

A boundary is *open* if it is an internal boundary of some base, otherwise it is *closed*. A section $\sigma = [i, \dots, i+k]$ is said to be *closed* if the boundaries i and $i+k$ are closed and all the boundaries between them are open.

Sometimes it will be convenient to subdivide all sections of Ω into *active* (denoted $A\Sigma_\Omega$) and *non-active* sections. For an item h denote by $\gamma(h)$ the number of bases containing h . An item h is called *free* if it meets no base, that is, if $\gamma(h) = 0$. Free variables are transported to the very end of the interval behind all items in Ω and they become non-active.

(D1) (*Closing a section*). Let σ be a section of Ω . The transformation (D1) makes the section σ closed. Namely, (D1) cuts all bases in Ω through the end-points of σ .

(D2) (*Transporting a closed section*). Let σ be a closed section of a generalized equation Ω . We cut σ out of the interval $[1, \rho_\Omega]$ together with all the bases on σ and put σ at the end of the interval or between any two consecutive closed sections of Ω . After that we correspondingly re-enumerate all the items and boundaries of the latter equation to bring it to the proper form. Clearly, the original equation Ω and the new one Ω' have the same solution sets and their coordinate groups are isomorphic.

(D3) (*Moving free variables to the right*). Suppose that Ω contains a free variable h_q in an active section. Here we close the section $[q, q+1]$ using (D1), transport it to the very end of the interval behind all items in Ω using (D2). In the resulting generalized equation Ω' the transported section becomes a non-active section.

(D4) (*Deleting a complete base*). A base μ of Ω is called *complete* if there exists a closed section σ in Ω such that $\sigma = [\alpha(\mu), \beta(\mu)]$.

Suppose μ is an active complete base of Ω and σ is a closed section such that $\sigma = [\alpha(\mu), \beta(\mu)]$. In this case using (ET5), we transfer all bases from μ to $\bar{\mu}$, then using (ET4) we remove the lone base μ together with the section σ .

(D5) (*Finding the kernel*). We will give a definition of eliminable base for an equation Ω that does not have any boundary connections. An active base $\mu \in A\Sigma_\Omega$ is called *eliminable* if at least one of the following holds

- (a) μ contains an item h_i with $\gamma(h_i) = 1$,
- (b) at least one of the boundaries $\alpha(\mu), \beta(\mu)$ is different from $1, \rho + 1$ and does not touch any other base (except for μ).

The process of finding the kernel works as follows. We cut the bases of Ω along all the boundary connections thus obtaining the equation without boundary connections, then consequently remove eliminable bases until no eliminable base is left in the equation. The resulting generalized equation is called the *kernel* of Ω and we denote it by $Ker(\Omega)$. It is easy to see that $Ker(\Omega)$ does not depend on a particular removal process. Indeed, if Ω has two different eliminable bases μ_1, μ_2 , and deletion of μ_i produces an equation Ω_i then by induction (on the number of eliminations) $Ker(\Omega_i)$ is uniquely defined for $i = 1, 2$. Obviously, μ_1 is still eliminable in Ω_2 , as well as μ_2 is eliminable in Ω_1 . Now eliminating μ_1 and μ_2 from Ω_2 and Ω_1 we get one and the same equation Ω_0 . By induction, $Ker(\Omega_1) = Ker(\Omega_0) = Ker(\Omega_2)$ hence the result. We say that an item h_i belongs to the kernel ($h_i \in Ker(\Omega)$), if h_i belongs to at least one base in the kernel. Notice that the kernel can be empty.

Also, for an equation Ω by $\overline{\Omega}$ we denote the equation which is obtained from Ω by deleting all free variables. Obviously,

$$G_{\Omega} = G_{\overline{\Omega}} * F(\overline{Y}),$$

where $\overline{Y}$ is the set of free variables in Ω .

Let us consider what happens in this process on the group level.

We start with the case when just one base is eliminated. Let μ be an eliminable base in $\Omega = \Omega(h_1, \dots, h_{\rho})$. Denote by Ω_1 the equation resulting from Ω by eliminating μ .

- (a) Suppose $h_i \in \mu$ and $\gamma(h_i) = 1$. Then the variable h_i occurs only once in Ω - precisely in the equation $s_{\mu} = 1$ corresponding to the base μ . Therefore, in the group G_{Ω} the relation $s_{\mu} = 1$ can be written as $h_i = w$, where w does not contain h_i . Using Tietze transformations we can rewrite the presentation of G_{Ω} as $G_{\Omega'}$, where Ω' is obtained from Ω by deleting s_{μ} and the item h_i . It follows immediately that

$$G_{\Omega_1} \simeq G_{\Omega'} * \langle h_i \rangle$$

and

$$G_{\Omega} \simeq G_{\Omega'} \simeq G_{\Omega_1} * F(B) \tag{1}$$

for some free group $F(B)$.

- (b) Suppose now that μ satisfies case (b) above with respect to a boundary i . Then in the equation $s_{\mu} = 1$ the variable h_{i-1} either occurs only once, or it occurs precisely twice and in this event the second occurrence of h_{i-1} (in $\overline{\mu}$) is a part of the subword $(h_{i-1}h_i)^{\pm 1}$. In both cases it is easy to see that the tuple

$$(h_1, \dots, h_{i-2}, s_{\mu}, h_{i-1}h_i, h_{i+1}, \dots, h_{\rho})$$

generates G . Therefore, by eliminating the relation $s_{\mu} = 1$ we can rewrite the presentation of G_{Ω} in the generators $\overline{Y} = (h_1, \dots, h_{i-2}, h_{i-1}h_i, h_{i+1}, \dots, h_{\rho})$. Observe also that any other equation $s_{\lambda} = 1$ ($\lambda \neq \mu$) of Ω either does not contain the variables h_{i-1}, h_i , or it contains them as parts of the subword $(h_{i-1}h_i)^{\pm 1}$, that is, any such a word s_{λ} can be expressed as a word $w_{\lambda}(\overline{Y})$ in terms of the generators $\overline{Y}$. This shows that

$$G_{\Omega} \simeq \langle \overline{Y} \mid \{w_{\lambda}(\overline{Y}) \mid \lambda \neq \mu\} \rangle \simeq G_{\Omega'},$$

where Ω' is a generalized equation obtained from Ω_1 by deleting the boundary i . Denote by Ω'' an equation obtained from Ω' by adding a free variable z to the right end of Ω' . It follows now that

$$G_{\Omega_1} \simeq G_{\Omega''} \simeq G_{\Omega} * \langle z \rangle$$

and

$$G_{\Omega} \simeq G_{\overline{\Omega'}} * F(K) \quad (2)$$

for some free group $F(K)$.

By induction on the number of steps in the process we obtain the following lemma.

Lemma 6. *If Ω is a generalized equation, then*

$$F_{\Omega} \simeq F_{\overline{Ker(\Omega)}} * F(K)$$

where $F(K)$ is a free group on K .

Proof. Let

$$\Omega = \Omega_0 \rightarrow \Omega_1 \rightarrow \cdots \rightarrow \Omega_l = Ker(\Omega)$$

be a linear elimination process. It is easy to see (by induction on l) that for every $j = 0, \dots, l-1$

$$\overline{Ker(\Omega_j)} = \overline{Ker(\overline{\Omega_j})}.$$

Moreover, if Ω_{j+1} is obtained from Ω_j as in the case (b) above, then (in the notations above)

$$\overline{Ker(\Omega_j)_1} = \overline{Ker(\Omega'_j)}.$$

Now the statement of the lemma follows from the remarks above and equalities (1) and (2). $\square$

- (D6) (*Linear elimination*). Suppose that in Ω there is h_i in an active section with $\gamma(h_i) = 1$ and such that $|h_i|$ is comparable with the length of the active section. In this case we say that Ω is *linear in h_i* .

If Ω is linear in h_i in an active section such that both boundaries i and $i+1$ are closed then we remove the closed section $[i, i+1]$ together with the lone base using (ET4).

If there is no such h_i but Ω is linear in some h_i in an active section such that one of the boundaries $i, i+1$ is open, say $i+1$, and the other is closed, then we perform (ET5) and μ -tie $i+1$ through the only base μ it intersects. Next, using (ET1) we cut μ in $i+1$ and then we delete the closed section $[i, i+1]$ by (ET4).

Suppose there is no h_i as above but Ω is linear in some h_i in an active section such that both boundaries i and $i+1$ are open. In addition, assume that there is a closed section σ containing exactly two (not matched) bases μ_1 and μ_2 , such that $\sigma = \sigma(\mu_1) = \sigma(\mu_2)$ and in the generalized equation $\tilde{\Omega}$ (see the derived transformation (D3)) all the bases obtained from μ_1, μ_2 by (ET1) in constructing $\tilde{\Omega}$ from Ω , do not belong to the kernel of $\tilde{\Omega}$. Here, using (ET5), we μ_1 -tie all the boundaries inside of μ_1 , then using (ET2) we transfer μ_2 onto $\overline{\mu_1}$, and remove μ_1 together with the closed section σ using (ET4).

Suppose now that Ω satisfies the first assumption of the previous paragraph and does not satisfy the second one. In this event we close the section $[i, i+1]$ using (D1) and remove it using (ET4).

Lemma 7. *Suppose that the process of linear elimination continues infinitely and there is a corresponding sequence of generalized equations*

$$\Omega \rightarrow \Omega_1 \rightarrow \cdots \rightarrow \Omega_k \rightarrow \cdots .$$

Then

- (a) ([23], Lemma 15) *The number of different generalized equations that appear in the process is finite. Therefore some generalized equation appears in this process infinitely many times.*
- (b) ([23], Lemma 15) *If $\Omega_j = \Omega_k$, $j < k$ then $\pi(j, k)$ is an isomorphism, invariant with respect to the kernel, namely $\pi(j, k)(h_i) = h_i$ for any variable h_i that belongs to some base in $\text{Ker}(\Omega)$.*
- (c) *The interval for the equation Ω_j can be divided into two disjoint parts, each being the union of closed sections, such that one part is a generalized equation $\text{Ker}(\Omega)$ and the other part is non-empty and corresponds to a generalized equation Ω' , such that $G_{\Omega'} = F(K)$ is a free group on variables K and $G_{\Omega} = G_{\text{Ker}(\Omega)} * F(K)$.*

Proof. We will only prove the third statement. In the process of linear elimination we do not remove or cut variables h_i that belong to the kernel. Since we do not cut variables that belong to the kernel, we can remove parts of bases that cover some of these variables only finitely many times. Therefore, bases that are cut in the process belong to the part of the interval that is disjoint from the $\text{Ker}(\Omega)$. Together with Lemma 6 this proves the third statement. $\square$

(D7) (*Tietze cleaning*). This transformation consists of four transformations performed consecutively

- (a) linear elimination: if the process of linear elimination goes infinitely we replace the equation by its kernel,
- (b) deleting all pairs of matched bases,
- (c) deleting all complete bases,
- (d) moving all free variables to the right.

(D8) (*Entire transformation*). We need a few definitions. A base μ of the equation Ω is called a *leading* base if $\alpha(\mu) = 1$. A leading base is said to be *maximal* (or a *carrier base*) if $\beta(\lambda) \leq \beta(\mu)$ for any other leading base λ . Let μ be a carrier base of Ω . Any active base $\lambda \neq \mu$ with $\beta(\lambda) \leq \beta(\mu)$ is called a *transfer* base (with respect to μ).

Suppose now that Ω is a generalized equation with $\gamma(h_i) \geq 2$ for each h_i in the active part of Ω and such that $|h_i|$ is comparable with the length of the active part. *Entire transformation* is a sequence of elementary transformations which are performed as follows. We fix a carrier base μ of Ω . We transfer all transfer bases from μ onto $\bar{\mu}$. Now, there exists some $i < \beta(\mu)$ such that $h_1, \dots, h_i$ belong to only one base μ , while h_{i+1} belongs to at least two bases. Applying (ET1) we cut μ along the boundary $i + 1$. Finally, applying (ET4) we delete the section $[1, i + 1]$.

4.3 Complexity of a generalized equation and Delzant-Potyagailo complexity $c(G)$ of a group G

Denote by ρ_A the number of variables h_i in all active sections of Ω , by $n_A = n_A(\Omega)$ the number of bases in all active sections of Ω , by ν' the number of open boundaries in the active sections, and by σ' the number of closed boundaries in the active sections.

For a closed section $\sigma \in \Sigma_\Omega$ denote by $n(\sigma), \rho(\sigma)$ the number of bases and, respectively, variables in σ .

$$\begin{aligned}\rho_A &= \rho_A(\Omega) = \sum_{\sigma \in A\Sigma_\Omega} \rho(\sigma), \\ n_A &= n_A(\Omega) = \sum_{\sigma \in A\Sigma_\Omega} n(\sigma).\end{aligned}$$

The *complexity* of the generalized equation Ω is the number

$$\tau = \tau(\Omega) = \sum_{\sigma \in A\Sigma_\Omega} \max\{0, n(\sigma) - 2\}.$$

Notice that the entire transformation (D5) as well as the cleaning process (D4) do not increase complexity of equations.

Below we recall Delzant-Potyagailo's result (see [12]). A family $\mathcal{C}$ of subgroups of a torsion-free group G is called *elementary* if

- (a) $\mathcal{C}$ is closed under taking subgroups and conjugation,
- (b) every $C \in \mathcal{C}$ is contained in a maximal subgroup $\overline{C} \in \mathcal{C}$,
- (c) every $C \in \mathcal{C}$ is small (does not contain F_2 as a subgroup),
- (d) all maximal subgroups from $\mathcal{C}$ are malnormal.

G admits a *hierarchy* over $\mathcal{C}$ if the process of decomposing G into an amalgamated product or an HNN-extension over a subgroup from $\mathcal{C}$, then decomposing factors of G into amalgamated products and/or HNN-extensions over a subgroup from $\mathcal{C}$ etc. eventually stops.

Proposition 1. ([12]) *If G is a finitely presented group without 2-torsion and $\mathcal{C}$ is a family of elementary subgroups of G then G admits a hierarchy over $\mathcal{C}$.*

Corollary 6. *If G is a finitely presented Λ -free group then G admits a hierarchy over the family of all abelian subgroups.*

There is a notion of complexity of a group G defined in [12] and denoted by $c(G)$. We will only use the following statement that follows from there.

Proposition 2. ([12]) *If G is a finitely presented Λ -free group. Let Γ be an abelian decomposition of G as a fundamental group of a graph of groups with more than one vertex and with each edge group being maximal abelian at least in one of its vertex groups. Then for each vertex group G_v , $c(G_v) < c(G)$.*

4.4 Rewriting process for Ω

In this section we describe a rewriting process (elimination process) for a generalized equation Ω corresponding to G . Performing the elimination process we eventually detect a decomposition of G as a free product or (if it is freely indecomposable) as the fundamental group of a graph of groups with vertex groups of three types: QH vertex groups, abelian vertex groups (corresponding to periodic structures, see below), non-QH, non-abelian vertex groups (we will call them *weakly rigid* meaning that we do not split them in this particular decomposition). We also can detect splitting of G as an HNN-extension with stable letter infinitely longer than generators of the abelian associated subgroups. After obtaining such a decomposition we continue the elimination process with the generalized equation corresponding to free factors of G or to weakly rigid subgroups of G (we will show that this generalized equation can be naturally obtained from the generalized equation

Ω . The Delzant-Potyagailo complexity of factors in a free decomposition and complexity of weakly rigid subgroups is smaller than the complexity of G (denoted $c(G)$) [12]. In the case of an HNN extension we will show that the complexity τ of the generalized equation corresponding to a weakly rigid subgroup is smaller than the complexity of Ω .

We assume that Ω is in standard form, namely, that transformations (ET3), (D3) and (D4) have been applied to Ω and that on each step we apply them to the generalized equation before applying any other transformation.

Let Ω be a generalized equation. We construct a path $T(\Omega)$ (with associated structures), as a directed path oriented from the root v_0 , starting at v_0 and proceeding by induction on the distance n from the root.

We start with a general description of the path $T(\Omega)$. For each vertex v in $T(\Omega)$ there exists a unique generalized equation Ω_v associated with v . The initial equation Ω is associated with the root v_0 , $\Omega_{v_0} = \Omega$. In addition there is a homogeneous system of linear equations Σ_v with integer coefficients on the lengths of variables of Ω_v . We take Σ_{v_0} to be empty. For each edge $v \rightarrow v'$ (here v and v' are the origin and the terminus of the edge) there exists an isomorphism $\pi(v, v') : G_{\Omega_v} \rightarrow G_{\Omega_{v'}}$ associated with $v \rightarrow v'$.

If

$$v_0 \rightarrow v_1 \rightarrow \cdots \rightarrow v_s \rightarrow u$$

is a dubpath of $T(\Omega)$, then by $\pi(v, u)$ we denote composition of corresponding isomorphisms

$$\pi(v, u) = \pi(v, v_1) \circ \cdots \circ \pi(v_s, u).$$

If $v \rightarrow v'$ is an edge then there exists a finite sequence of elementary or derived transformations from Ω_v to $\Omega_{v'}$ and the isomorphism $\pi(v, v')$ is a composition of the isomorphisms corresponding to these transformations. We also assume that active (and non-active) sections in $\Omega_{v'}$ are naturally inherited from Ω_v , if not said otherwise. Recall that initially all sections are active.

Suppose the path $T(\Omega)$ is constructed by induction up to a level n and suppose v is a vertex at distance n from the root v_0 . We describe now how to extend the path from v . The construction of the outgoing edge at v depends on which case described below takes place at the vertex v . There are three possible cases.

- **Linear case:** there exists h_i in the active part such that $|h_i|$ is comparable with the length of the active part and $\gamma(h_i) = 1$.
- **Quadratic and almost quadratic case:** $\gamma(h_i) = 2$ for all h_i in the active part such that $|h_i|$ is comparable with the length of the active part.
- **General JSJ case:** $\gamma(h_i) \geq 2$ for all h_i in the active part such that $|h_i|$ is comparable with the length of the active part, and there exists such h_i that $\gamma(h_i) > 2$.

4.4.1 Linear case

We apply Tietze cleaning (D7) at the vertex v_n if it is possible. We re-write the system of linear equations Σ_{v_n} in new variables and obtain a new system $\Sigma_{v_{n+1}}$.

If $\Omega_{v_{n+1}}$ splits into two parts, $\Omega_{v_{n+1}}^{(1)} = \text{Ker}(\Omega_{v_n})$ and $\Omega_{v_{n+1}}^{(2)}$ that corresponds to a free group $F(K)$, then we put the free group section $\Omega_{v_{n+1}}^{(2)}$ into a non-active part and thus decrease the complexity. It may happen that the kernel is empty, then the process terminates.

If it is impossible to apply Tietze cleaning (that is $\gamma(h_i) \geq 2$ for any h_i in the active part of Ω_v comparable to the length of the active part), we apply the entire transformation.

Termination condition: Ω_v does not contain active sections. In this case the vertex v is called a *leaf* or an *end vertex*.

4.4.2 Quadratic case

Suppose Ω_v satisfies the condition $\gamma_i \leq 2$ for each h_i in the active part and $\gamma_i = 2$ for each h_i in the active part comparable with the length of the active part. First of all, we fill in all the h'_i s in the active part such that $\gamma_i = 1$ by new (infinitely short) bases μ such that $\bar{\mu}$ covers a new variable that we add to the non-active part.

We apply the entire transformation (D8), then apply Tietze cleaning (D7), if possible, then again apply entire transformation, etc. In this process we, maybe, will remove some pairs of matching bases decreasing the complexity. Eventually we either end up with empty active part or the process will continue infinitely, and the number of bases in the active part will be constant. Let $\alpha_0 + 1$ be the leftmost boundary that belongs to a base not participating in the process.

Let us analyze the structure of G_{Ω_v} in this case. Let σ be a section in the active part $\sigma = [h_1, \dots, h_{\alpha_0}]$. If σ is not closed, then cutting the second base at α_0 we make σ a closed section. Let F_1 be a free group with basis $\{h_1, \dots, h_{\alpha_0}\}$. If σ contains an open boundary j then we can consider a new generalized equation Ω'_v obtained from Ω_v by replacing the product $h_{j-1}h_j$ by a new variable h'_{j-1} and represent $G_{\Omega_v} = G_{\Omega'_v} * \langle h_j \rangle$. Therefore we can suppose that σ does not contain open boundaries. We say that μ is a *quadratic* base if σ contains μ and $\bar{\mu}$, otherwise μ is a *quadratic-coefficient base*. Denote the set of quadratic-coefficient bases by C . Suppose that σ contains quadratic bases. Let F_1/σ be the quotient of F_1 over the normal closure of elements $h[\alpha(\mu), \beta(\mu)]h[\alpha(\bar{\mu}), \beta(\bar{\mu})]^{-1}$. Let $\mathcal{M}$ be a set that contains exactly one representative of each pair of double bases on σ and contains also each base μ such that $\mu \in \sigma$ and $\bar{\mu} \notin \sigma$. If we identify each base on σ with its double then the product $h_1 \cdots h_i$ can be written as a product of bases from $\mathcal{M}$ in exactly two different ways: $\mu_{i_1} \cdots \mu_{i_k}$ and $\mu_{j_1} \cdots \mu_{j_t}$. Then F_1/σ is isomorphic to the quotient of the free group $F(\mathcal{M})$ over the relation $\mu_{i_1} \cdots \mu_{i_k} = \mu_{j_1} \cdots \mu_{j_t}$. Every element of $\mathcal{M}$ occurs in this relation at most twice. Applying an automorphism of $F(\mathcal{M})$ identical on C , we can obtain another basis $X \cup T \cup C$ of this group such that in this basis the relation has form of a standard quadratic equation in variables from X with coefficients in $F(C)$ (see [29], Section 1.7). Note that the variables from T do not participate in any relations. The quadratic equation corresponding to σ can be written in the standard form with coefficients expressed in terms of non-active variables.

Here we have to give some definitions.

Definition 6. A standard quadratic equation over the group G is an equation of the one of the following forms (below d, c_i are nontrivial elements from G):

$$\prod_{i=1}^n [x_i, y_i] \prod_{i=1}^m z_i^{-1} c_i z_i d = 1, \quad n, m \geq 0, m + n \geq 1; \quad (3)$$

$$\prod_{i=1}^n x_i^2 \prod_{i=1}^m z_i^{-1} c_i z_i d = 1, \quad n, m \geq 0, n + m \geq 1, \quad (4)$$

where $c_1, \dots, c_m, d$ are coefficients.

Let W be a strictly quadratic word over a group G . Then there is a G -automorphism $f \in \text{Aut}_G(G[X])$ such that W^f is a standard quadratic word over G .

To each quadratic equation one can associate a punctured surface. For example, the orientable surface associated to equation 3 will have genus n and $m + 1$ punctures.

Put

$$\kappa(S) = |X| + 1,$$

Definition 7. Let $S = 1$ be a standard quadratic equation over a group G which has a solution in G . The equation $S(X) = 1$ is regular if $\kappa(S) \geq 4$ (equivalently, the Euler characteristic of the corresponding punctured surface is at most -2) and there is a non-commutative solution of $S(X) = 1$ in G , or it is an equation of the type $[x, y]d = 1$, or $[x_1, y_1][x_2, y_2] = 1$.

If the quadratic equation corresponding to σ is regular, then G_{Ω_v} has a presentation as a fundamental group of a graph of groups with QH-subgroup corresponding to this equation. (For example, the QH-subgroup corresponding to the standard equation (3) has the presentation

$$\langle x_1, y_1, \dots, x_n, y_n, p_1, \dots, p_{m+1} \mid \prod_{i=1}^n [x_i, y_i] \prod_{i=1}^{m+1} p_i = 1 \rangle,$$

and the edge groups are $\langle p_i \rangle, i = 1, \dots, m+1$.)

If the equation is not regular and there are quadratic bases, then G_{Ω_v} splits as an HNN-extension over an abelian subgroup. If there are infinitely small h_i 's in the quadratic part such that $\gamma(h_i) = 1$ then G_{Ω_v} splits as a free product of the group of the generalized equation on the non-quadratic part (obtained by removing the quadratic part and all bases that have doubles in quadratic part) and a free group.

We say that a base *participates* in the infinite process of entire transformation if it participates infinitely many times as a leading or transfer base.

Lemma 8. *If σ has quadratic-coefficient bases, and the entire transformation goes infinitely, then after a finite number of steps there will be quadratic bases that have length infinitely larger than all participating quadratic coefficient bases.*

Proof. After finite number of steps quadratic-coefficient bases can only be used as transfer bases (otherwise quadratic bases would be leaving σ , and this cannot happen infinitely many times). Every time such a base of length ℓ used as a transfer base, the length of the quadratic part decreases by ℓ . For the process to go infinitely, ℓ must be infinitely less than the length of the quadratic part. Therefore there is a quadratic base that have length infinitely larger than all participating quadratic coefficient bases. $\square$

If σ does not have quadratic-coefficient bases then $G_{R(\Omega_v)}$ splits as a free product with one factor being a closed surface group or a free group.

We move σ into a non-active part and thus decrease the complexity τ . Eventually in this case we will make the active part empty.

4.4.3 Almost Quadratic case

In this case $\gamma_i = 2$ for all h_i in the active part such that $|h_i|$ is comparable with the length of the active part (some short bases may be covered once or more than twice). This case is, actually, a part of the case considered in the next subsection, but we consider it separately to make the subdivision into the cases depend only on γ_i for the variables h_i of the maximal height.

We apply the transformation (D1) to close quadratic sections that begin with bases of the maximal height and put them in front of the interval. This may increase the complexity by two times the number of closed sections that we put in front of the interval applying (D1). Then we apply the entire transformation. Every time when the content of one of the closed sections in front of the interval is moved to the other closed section, the complexity decreases by two. If the process goes infinitely then eventually the number of quadratic bases of maximal height does not decrease. After a finite number of steps participating quadratic-coefficient bases can only be used as transfer bases. And, therefore, there will be participating quadratic bases that have length infinitely larger than all participating quadratic coefficient bases.

We fill in all the h_i 's in the active part such that $\gamma_i = 1$ by new (infinitely short) bases μ such that $\bar{\mu}$ covers a new free variable in the non-active part appearing only once. Then we work with the quadratic part the same way as in the quadratic case. We repeat the described transformation until there is no quadratic base on the active part that has length comparable with the length of the remaining active part. Then we consider the remaining generalized equation in the active

part. We remove from the active part doubles of all quadratic coefficient bases that belong to non-active part (doing this we may create new boundaries). We will, certainly remember the relations corresponding to these pairs of bases. In this case the remaining generalized equation has smaller complexity. Relations corresponding to the quadratic sections that we made non-active show that G_Ω is an HNN-extension of the subgroup generated by the variables in the active part and (maybe) a free group.

Removing these double bases we have to add equations to $\Sigma_{v_{i+1}}$ that guarantee that the associated cyclic subgroups are generated by elements of the same length.

4.4.4 General JSJ-case

Generalized equation Ω_v satisfies the condition $\gamma_i \geq 2$ for each h_i in the active part such that $|h_i|$ is comparable with the length of the active part, and $\gamma_i > 2$ for at least one h_i . First of all, we fill in all the h_i 's in the active part such that $\gamma_i = 1$ by new (infinitely short) bases with doubles corresponding to free variables in the non-active part. We apply the transformation (D1) to close the quadratic part and put it in front of the interval.

(a) QH-subgroup case. Suppose that the entire transformation of the quadratic part (D8) goes infinitely. Then the quadratic part of Ω_v (or the initial section from the beginning of the quadratic part until the first base on the quadratic part that does not participate in the entire transformation) corresponds to a QH-vertex or to the representation of G_Ω as an HNN-extension, and there is a quadratic base (on this section) that is infinitely longer than all the quadratic coefficient bases (on this section). We work with the quadratic part the same way as in the quadratic case until there is no quadratic base satisfying the condition above. Then we make the quadratic section non-active, and consider the remaining generalized equation where we remove doubles of all the quadratic coefficient bases. We certainly have to remember that the bases that we removed express some variables in the quadratic part (that became non-active) in the variables in the active part. We have to add an equation to $\Sigma_{v_{i+1}}$ that guarantees that the associated cyclic subgroups are generated by elements of the same length. In this case the subgroup of G_Ω that is isomorphic to the coordinate group of the new generalized equation in active part is a vertex group in an abelian splitting of G_Ω and has smaller Delzant–Potyagailo complexity.

(b) QH-shortening

Lemma 9. *Suppose that the quadratic part of Ω does not correspond to the HNN-splitting of G_Ω (there are only quadratic coefficient bases), or, we cannot apply the entire transformation to the quadratic part infinitely. In this case either G_Ω is a non-trivial free product or, applying the automorphism of G_Ω , one can replace the words corresponding to the quadratic bases in the quadratic part by such words that in the new solution H^+ of Ω the length of the quadratic part is bounded by some function $f_1(\Omega)$ times the length of the non-quadratic part. Solution H^+ can be chosen consistent with H .*

Proof. We apply entire transformation to the quadratic part. This corresponds to a sequence of Tietze transformations of generators of G_Ω . Since the entire transformation does not go infinitely, some of the bases will eventually be transferred to the non-quadratic part (therefore, their length is bounded by the length of the non-quadratic part). If not all of them are transferred, then reincarnations of some of the other participating bases must form matching pairs giving a representation of G_Ω as a free product. $\square$

If there is a matching pair, we replace G_Ω by the group obtained by removing a cyclic free factor corresponding to a matching pair. We also replace Ω by the generalized equation obtained by removing the matching pair. The Delzant–Potyagailo complexity decreases.

(c) Abelian splitting: short shift.

Proposition 3. *Suppose Ω_v satisfies the following condition: the carrier base μ of the equation Ω_v intersects with its dual $\bar{\mu}$ (form an overlapping pair) and is at least twice longer than $|\alpha(\bar{\mu}) - \alpha(\mu)|$. Then G_{Ω_v} either splits as a fundamental group of a graph of groups that has a free abelian vertex group or splits as an HNN-extension with abelian associated subgroups.*

The proof uses the technique of so-called periodic structures introduced by Razborov in his Ph.D thesis and almost repeats the proof given in [23] to show that the coordinate group of a generalized equation splits in this case as a fundamental group of a graph of groups that has a free abelian vertex group or splits as an HNN-extension with abelian associated subgroups. Nevertheless, we have to adjust the proof to the case of infinite non-Archimedean words. For convenience of the reader we give a detailed proof of the proposition.

We construct an auxiliary equation $\widehat{\Omega}_v$ (which does not occur in $T(\Omega)$) as follows. Firstly, we add a new constant non-active section $[\rho_v + 1, \rho_v + 2]$ to the right of all sections in Ω_v (in particular, h_{ρ_v+1} is a new free variable). Secondly, we introduce a new pair of bases $(\lambda, \bar{\lambda})$ such that

$$\alpha(\lambda) = 1, \beta(\lambda) = \beta(\bar{\mu}), \alpha(\bar{\lambda}) = \rho_v + 1, \beta(\bar{\lambda}) = \rho_v + 2.$$

Notice that Ω_v can be obtained from $\widehat{\Omega}_v$ by (ET4): deleting $\bar{\lambda}$ together with the closed section $[\rho_v + 1, \rho_v + 2]$. Let

$$\widehat{\pi}_v : G_{\Omega_v} \rightarrow G_{\widehat{\Omega}_v}$$

be the isomorphism induced by (ET4). General JSJ-case still holds for $\widehat{\Omega}_v$, but now λ is the carrier base. Applying to $\widehat{\Omega}_v$ the entire transformation we obtain a new vertex $\Omega_{v'}$ together with the isomorphism

$$\eta_{v'} : G_{\widehat{\Omega}_v} \rightarrow G_{\Omega_{v'}}.$$

As a culmination of the above process of adding auxiliary edges we get a generalized equation Ω' , which consists of finitely many closed sections $\sigma_1, \dots, \sigma_p$ such that there exists an overlapping pair $(\mu_i, \bar{\mu}_i)$ for each σ_i , $i \in [1, p]$.

Let $\sigma = \sigma_{i_0}$, where $i_0 \in [1, p]$ have the overlapping pair $(\mu, \bar{\mu})$ such that

$$|[\alpha(\mu), \alpha(\bar{\mu})]| = \max_{i \in [1, p]} |[\alpha(\mu_i), \alpha(\bar{\mu}_i)]|.$$

Without loss of generality we can assume $i_0 = 1$, that is, $\sigma = [1, \beta(\bar{\mu})]$.

A *periodic structure* on Ω' is a pair $\langle \mathcal{P}, R \rangle$, where

1. $\mathcal{P}$ is a set consisting of some items h_i , some bases λ , and some closed sections σ from Ω' such that the following conditions are satisfied:
 - (a) if $h_i \in \mathcal{P}$ and $h_i \in \lambda$ then $\lambda \in \mathcal{P}$,
 - (b) if $\lambda \in \mathcal{P}$, then $\bar{\lambda} \in \mathcal{P}$,
 - (c) if $\lambda \in \mathcal{P}$ and $\lambda \in \sigma$, then $\sigma \in \mathcal{P}$,
 - (d) there exists a function $\mathcal{X}$ mapping the set of closed sections from $\mathcal{P}$ into $\{-1, +1\}$ such that for every $\lambda, \sigma_1, \sigma_2 \in \mathcal{P}$, the condition that $\lambda \in \sigma_1$ and $\bar{\lambda} \in \sigma_2$ implies $\varepsilon(\lambda) \cdot \varepsilon(\bar{\lambda}) = \mathcal{X}(\sigma_1) \cdot \mathcal{X}(\sigma_2)$.
2. R is an equivalence relation on the set of boundaries $\mathcal{B}$ belonging to closed sections from $\mathcal{P}$ (if a boundary l belongs to two closed section $\sigma_{left}(l) = [i, l], \sigma_{right}(l) = [l, j]$ from $\mathcal{P}$ then instead of l we add to $\mathcal{B}$ formal copies l_{left}, l_{right} of l , which are boundaries of $\sigma_{left}(l)$ and $\sigma_{right}(l)$ respectively), defined as follows: if $\lambda \in \mathcal{P}$ then

$$\alpha(\lambda) \sim_R \alpha(\bar{\lambda}), \beta(\lambda) \sim_R \beta(\bar{\lambda}) \text{ if } \varepsilon(\lambda) = \varepsilon(\bar{\lambda})$$

$$\alpha(\lambda) \sim_R \beta(\bar{\lambda}), \beta(\lambda) \sim_R \alpha(\bar{\lambda}) \text{ if } \varepsilon(\lambda) = -\varepsilon(\bar{\lambda}).$$

We call items, bases and sections *long* if they belong to $\mathcal{P}$ (the rest items, bases and sections of Ω' are called *short* respectively).

A *period* is just a cyclically reduced Λ -word u . A reduced infinite word w is called *u -periodic* for a period u if $w^{-1} * u * w = v$, where $|u| = |v|$. Since u is cyclically reduced it follows that either $w^{-1} * u = w^{-1} \circ u$, or $u * w = u \circ w$. We are going to distinguish two cases of periodicity:

1. w is *unbounded u -periodic* if $ht(w) \geq ht(u)$,
2. w is *bounded u -periodic* if $ht(w) = ht(u)$ and $w = u^k \circ u_1$, where $k \geq 2$ and $u = u_1 \circ u_2$.

Observe that in both cases w begins with a power of u , but in the former case the power is unbounded, while in the latter case it is bounded (hence the terms used).

Lemma 10. *Let $w_1, w_2 \in CDR(\Lambda, Z)$, where $|w_1| \geq |w_2|$, be u -periodic for some period $u \in CDR(\Lambda, Z)$. Assume that $w_1^{-1} * u * w_1 = v_1$, $w_2^{-1} * u * w_2 = v_2$. If $w_1^{-1} * w_2$ is defined in $CDR(\Lambda, Z)$ then w_2 cancels completely in this product. Moreover, $w_1^{-1} * w_2$ is v_1^{-1} -periodic.*

Proof. Follows from the definition of u -periodic words. □

Lemma 11. *If $w \in CDR(\Lambda, Z)$ is both u - and v -periodic for some periods $u, v \in CDR(\Lambda, Z)$ then $[u, v] = 1$. In particular, u is unbounded v -periodic if $ht(u) > ht(v)$.*

Proof. If $ht(u) = ht(v)$ then commutativity follows from Lemma 3.3 [22] since powers of u and v have a common initial segment of length greater than $|u| + |v|$. If $ht(u) > ht(v)$ then from v -periodicity of w it follows that u is v -periodic. Now, since w contains any natural power of u as an initial subword it follows that the terminal subword of u of length $|v|$ must coincide with v and commutativity follows. □

Let U be a solution of the equation Ω' , where $h_i \rightarrow U_i$. According to our assumption, X_μ is P -periodic, where $P = U[1, \alpha(\bar{\mu})]$. Since μ and $\bar{\mu}$ are products of items then every U_i , $i \in [\alpha(\bar{\mu}), \beta(\mu)]$, such that $|U_i| \geq 2|P|$, is Q_i -periodic, where Q_i is a Λ -word such that $|Q_i| = |P|$. Moreover, we can associate with U a periodic structure $\mathcal{P}(U, P) = \langle \mathcal{P}, R \rangle$ as follows. We construct $\mathcal{P}$ as a union of the chain of sets

$$\mathcal{P}_1 \subseteq \mathcal{P}_2 \subseteq \dots \subseteq \mathcal{P}_k,$$

where $k \in \mathbb{N}$. To construct $\mathcal{P}_0$ we add to it σ_1 and all Q -periodic items $h \in \sigma_1$, where $|Q| = |P|$, together with any base λ (and its dual) containing h . Now, assume that $\mathcal{P}_i$ is constructed. If there exists a base $\lambda \in \mathcal{P}_i$ such that $\bar{\lambda} \in \sigma_j$ then we set

$$\mathcal{P}_{i+1} = \mathcal{P}_i \cup \{\sigma_j\} \cup \{h, \lambda \in \sigma_j \mid h \text{ is a } Q\text{-periodic item where } |Q| = |P|, \lambda \text{ is a base containing } h\}.$$

Since Ω' contains only finitely many closed sections then this process stops.

Lemma 12. *Let U be a solution of the equation Ω' and $\mathcal{P}(U, P)$ be a periodic structure corresponding to U . Then every base (item) $\lambda \in \mathcal{P}$ is Q_λ -periodic, where $|Q_\lambda| = |P|$ and Q_λ is conjugate to P in G .*

Proof. It is enough to prove the statement for bases - the argument for items is similar. We are going to use the induction on k , where k is the number of sections in $\mathcal{P}$

By our assumption (short shift) we have $\mu \in \mathcal{P}$. It follows that any base $\lambda \in \sigma_1$, such that $| \lambda | \geq 2|P|$, is Q_λ -periodic, where $|Q_\lambda| = |P|$. Moreover, if $w = U[\alpha(\bar{\mu}), \alpha\lambda]$ then $w \in G$ and $w^{-1} * P * w = Q_\lambda$. Hence, all bases from $\mathcal{P}_1$ have the properties stated in the lemma.

Suppose we have the statement of the lemma for all bases from $\mathcal{P}_{k-1}$. By definition, $\mathcal{P}_k$ is obtained from $\mathcal{P}_{k-1}$ by adding a section σ_k , such that there exists $\lambda \in \mathcal{P}_{k-1}$ and $\bar{\lambda} \in \sigma_k$, as well as all appropriate items and bases of σ_k . Observe that since μ_k is Q -periodic, where $Q = U[\alpha(\mu_k), \alpha(\bar{\mu}_k)]$,

it follows that $\bar{\lambda}$ is Q' -periodic, where $|Q'| = |Q|$ and Q' is conjugate to Q by means of $g \in G$. At the same time, by the induction hypothesis, λ is Q_λ -periodic, where $|Q_\lambda| = |P|$. Hence, by Lemma 11 we have $[Q', Q_\lambda] = 1$ and it follows that $[Q, g^{-1} * Q_\lambda * g] = 1$, where $|g^{-1} * Q_\lambda * g| = |P|$. Now, since $|P| \geq |Q|$ and $|\mu_k| \geq |\lambda|$ it follows that μ_k is $g^{-1} * Q_\lambda * g$ -periodic. The statement of the lemma for all bases in σ_k of the length at least $2|P|$ follows automatically now. $\square$

Remark 4. From the construction of $\mathcal{P}(U, P)$ and Lemma 12 it follows that if $\sigma_i \in \mathcal{P}$ with an overlapping pair $(\mu_i, \bar{\mu}_i)$ and if $l \in [\alpha(\sigma_i), \beta(\sigma_i)]$ then there exists a period Q_l conjugate in G either to P or P^{-1} such that

- (i) if $ht([\alpha(\sigma_i), l]) > ht(P)$ then $U[\alpha(\sigma_i), l]$ has any natural power of Q_l as a terminal subword,
- (ii) if $ht([l, \beta(\sigma_i)]) > ht(P)$ then $U[l, \beta(\sigma_i)]$ has any natural power of Q_l as an initial subword,
- (iii) if $ht([\alpha(\sigma_i), l]) \leq ht(P)$ then $U[\alpha(\sigma_i), l] = Q'' \circ Q_l^{k_l}$, where $Q_l = Q' \circ Q''$,
- (iv) if $ht([l, \beta(\sigma_i)]) \leq ht(P)$ then $U[l, \beta(\sigma_i)] = Q_l^{k_l} \circ Q'$, where $Q_l = Q' \circ Q''$.

Lemma 13. $\mathcal{P}(U, P)$ corresponding to a solution U of Ω' is a periodic structure on Ω' .

Proof. Parts 1(a), 1(b) and 1(c) of the definition of periodic structure hold by construction.

Set $\mathcal{X}(\sigma_1) = 1$. If there is a base $\lambda \in \sigma_1$ such that $\bar{\lambda} \in \sigma_2$ then we set $\mathcal{X}(\sigma_2) = \varepsilon(\lambda)\varepsilon(\bar{\lambda})$. Observe that the sign of $\mathcal{X}(\sigma_2)$ is well-defined since if there exists another base $\lambda_1 \in \sigma_1$ such that $\bar{\lambda}_1 \in \sigma_2$ and $\varepsilon(\lambda)\varepsilon(\bar{\lambda}) \neq \varepsilon(\lambda_1)\varepsilon(\bar{\lambda}_1)$ then it follows that there exists a group element conjugating P into P^{-1} , which is impossible. Hence 1(d) holds.

By Remark 4 if two boundaries l_1, l_2 belong to sections from $\mathcal{P}$ then there exist Q_{l_1}, Q_{l_2} conjugate in G either to P or P^{-1} such that some of the conditions listed in the remark hold. Thus, we define a relation R as follows: $l_1 \sim_R l_2$ if and only if $Q_{l_1} = Q_{l_2}$, and part 2 follows immediately. $\square$

Now let us fix a nonempty periodic structure $\langle \mathcal{P}, R \rangle$. Item (d) allows us to assume (after replacing the variables $h_i, \dots, h_{j-1}$ by $h_{j-1}^{-1}, \dots, h_i^{-1}$ on those sections $[i, j] \in \mathcal{P}$ for which $\mathcal{X}([i, j]) = -1$) that $\varepsilon(\mu) = 1$ for all $\mu \in \mathcal{P}$. For a boundary k , we will denote by (k) the equivalence class of the relation R to which it belongs.

Let us construct an oriented graph Γ whose set of vertices is the set of R -equivalence classes. For each long h_k we introduce an oriented edge e leading from (k) to $(k+1)$ and an inverse edge e^{-1} leading from $(k+1)$ to (k) . This edge e is assigned the label $h(e) = h_k$ (respectively, $h(e^{-1}) = h_k^{-1}$). For every path $r = e_1^{\pm 1} \cdots e_s^{\pm 1}$ in the graph Γ , we denote by $h(r)$ its label $h(e_1^{\pm 1}) \cdots h(e_s^{\pm 1})$.

The periodic structure $\langle \mathcal{P}, R \rangle$ is called *connected*, if the graph Γ is connected. Suppose first that $\langle \mathcal{P}, R \rangle$ is connected. We can also suppose that each boundary of Ω' is a boundary between two bases.

Lemma 14. Let U be a solution of Ω' and $\langle \mathcal{P}, R \rangle = \mathcal{P}(U, P)$. If p is a path in the graph Γ from (i) to (j) then $U[p]^{-1} * Q_i * U[p] = Q_j$ in G .

Proof. If $p = h_{i_1} \cdots h_{i_k}$ then all products between U_j , $j \in \{i_1, \dots, i_k\}$ are defined and the required result follows from Lemma 10. $\square$

Lemma 15. Let U be a solution of Ω' and $\langle \mathcal{P}, R \rangle = \mathcal{P}(U, P)$. If c_1, c_2 are cycles in the graph Γ at the vertex $v = (k)$ then $[U[c_1], U[c_2]] = 1$ in G .

Proof. Assume at first that $(k) = (1)$. There exists a loop c at (1) such that $U[c] = P$. By Lemma 14 $U[p]^{-1} * P * U[p] = P$ in G for every path p from (1) to (1) , so $[U[c_1], P] = [U[c_2], P] = 1$ and from the CSA-property of G the required follows.

If $(k) \neq (1)$ then, since Γ is connected it follows that there exists a path p from v to v_0 . Moreover, by Lemma 14 we have $U[p]^{-1} * Q_k * U[p] = P$ in G . So, by the CSA-property commutativity of cycles at (k) follows from commutativity of cycles at (1) which is already proved. $\square$

Ω' is called *periodized* with respect to a given periodic structure $\langle \mathcal{P}, R \rangle$ if for every two cycles c_1 and c_2 with the same initial vertex in the graph Γ there is a relation $[h(c_1), h(c_2)] = 1$ in $G_{\Omega'}$. Observe that according to Lemma 15 Ω' is periodized with respect to $\mathcal{P}(U, P)$ for any solution U .

Let Γ_0 be the subgraph of the graph Γ having the same set of vertices and consisting of the edges e whose labels do not belong to $\mathcal{P}$. Choose a maximal sub-forest T_0 in the graph Γ_0 and extend it to a maximal sub-forest T of the graph Γ . Since $\langle \mathcal{P}, R \rangle$ is connected by assumption, it follows that T is a tree. Let v_0 be an arbitrary vertex of the graph Γ and $r(v_0, v)$ the (unique) path from v_0 to v all of whose vertices belong to T . For every edge $e = (v, v')$ not lying in T , we introduce a cycle $c_e = r(v_0, v)e(r(v_0, v'))^{-1}$. Then the fundamental group $\pi_1(\Gamma, v_0)$ is generated by the cycles c_e (see, for example, the proof of Proposition 3.2.1 [29]).

Furthermore, the set of elements

$$\{h(e) \mid e \in T\} \cup \{h(c_e) \mid e \notin T\} \quad (5)$$

forms a basis of the free group with the set of generators $\{h_k \mid h_k \in \sigma\}$. If $\lambda \in \mathcal{P}$, then $(\beta(\lambda)) = (\beta(\bar{\lambda}))$, $(\alpha(\lambda)) = (\alpha(\bar{\lambda}))$ by the definition of R and, consequently, the word

$$h[\alpha(\lambda), \beta(\lambda)] h[\alpha(\bar{\lambda}), \beta(\bar{\lambda})]^{-1}$$

is the label of a cycle $c'(\lambda)$ from $\pi_1(\Gamma, (\alpha(\lambda)))$. Let $c(\lambda) = r(v_0, (\alpha(\lambda))) c'(\lambda) r(v_0, (\alpha(\lambda)))^{-1}$. Then

$$h(c(\lambda)) = u h[\alpha(\lambda), \beta(\lambda)] h[\alpha(\bar{\lambda}), \beta(\bar{\lambda})]^{-1} u^{-1}, \quad (6)$$

where u is a certain word. Since $c(\lambda) \in \pi_1(\Gamma, v_0)$, it follows that $c(\lambda) = b_\lambda(\{c_e \mid e \notin T\})$, where b_λ is a certain word in the indicated generators which can be effectively constructed (see Proposition 3.2.1 [29]).

Let $\tilde{b}_\lambda$ denote the image of the word b_λ in the abelianization of $\pi(\Gamma, v_0)$. Denote by $\tilde{Z}$ the free abelian group consisting of formal linear combinations $\sum_{e \notin T} n_e \tilde{c}_e$ ($n_e \in \mathbb{Z}$), and by $\tilde{B}$ its subgroup generated by the elements $\tilde{b}_\lambda$ ($\lambda \in \mathcal{P}$) and the elements $\tilde{c}_e$ ($e \notin T, h(e) \notin \mathcal{P}$). Let $\tilde{A} = \tilde{Z}/\tilde{B}$, $T(\tilde{A})$ the torsion subgroups of the group $\tilde{A}$, and $\tilde{Z}_1$ the preimage of $T(\tilde{A})$ in $\tilde{Z}$. The group $\tilde{Z}/\tilde{Z}_1$ is free, therefore, there exists a decomposition of the form

$$\tilde{Z} = \tilde{Z}_1 \oplus \tilde{Z}_2, \quad \tilde{B} \subseteq \tilde{Z}_1, \quad |\tilde{Z}_1 : \tilde{B}| < \infty. \quad (7)$$

Note that it is possible to express effectively a certain basis $\tilde{c}^{(1)}, \tilde{c}^{(2)}$ of the group $\tilde{Z}$ in terms of the generators $\tilde{c}_e$ so that for the subgroups $\tilde{Z}_1$ and $\tilde{Z}_2$ generated respectively by $\tilde{c}^{(1)}$ and $\tilde{c}^{(2)}$, the relation (7) holds. It suffices, for instance, to look through the bases one by one, using the fact that under the condition $\tilde{Z} = \tilde{Z}_1 \oplus \tilde{Z}_2$ the relations $\tilde{B} \subseteq \tilde{Z}_1$, $|\tilde{Z}_1 : \tilde{B}| < \infty$ hold if and only if the generators of the groups $\tilde{B}$ and $\tilde{Z}_1$ generate the same linear subspace over $\mathbf{Q}$, and the latter is easily verified algorithmically. Notice, that a more economical algorithm can be constructed by analyzing the proof of the classification theorem for finitely generated abelian groups. By Proposition 1.4.4 [29], one can effectively construct a basis $\tilde{c}^{(1)}, \tilde{c}^{(2)}$ of the free (non-abelian) group $\pi_1(\Gamma, v_0)$ so that $\tilde{c}^{(1)}, \tilde{c}^{(2)}$ are the natural images of the elements $\tilde{c}^{(1)}, \tilde{c}^{(2)}$ in $\tilde{Z}$.

Now assume that $\langle \mathcal{P}, R \rangle$ is an arbitrary periodic structure on Ω' , not necessarily connected. Let $\Gamma_1, \dots, \Gamma_r$ be the connected components of the graph Γ . The labels of edges of the component Γ_i form in Ω' a union of closed sections from $\mathcal{P}$. Moreover, if a base $\lambda \in \mathcal{P}$ belongs to such a section, then its dual $\bar{\lambda}$ also does by definition of periodic structure. Therefore, by taking for $\mathcal{P}_i$ the set of labels of edges from Γ_i belonging to $\mathcal{P}$, sections to which these labels belong, and bases $\lambda \in \mathcal{P}$ belonging to these sections, and restricting in the corresponding way the relation R , we obtain a periodic connected structure $\langle \mathcal{P}_i, R_i \rangle$ with the graph Γ_i .

The notation $\langle \mathcal{P}', R' \rangle \subseteq \langle \mathcal{P}, R \rangle$ means that $\mathcal{P}' \subseteq \mathcal{P}$ and the relation R' is a restriction of the relation R . In particular, $\langle \mathcal{P}_i, R_i \rangle \subseteq \langle \mathcal{P}, R \rangle$ in the situation described in the previous paragraph. Since Ω' is periodized, the periodic structure must be connected.

Let $e_1, \dots, e_m$ be all the edges of the graph Γ from $T - T_0$. Since T_0 is the spanning forest of the graph Γ_0 , it follows that $h(e_1), \dots, h(e_m) \in \mathcal{P}$. Let $F(\Omega')$ be a free group generated by the variables of Ω' . Consider in the group $F(\Omega')$ a new basis

$$Y = \{\bar{t}, \{h(e) \mid e \in T\}, h(\bar{c}^{(1)}), h(\bar{c}^{(2)})\},$$

where variables $\bar{t}$ do not belong to the closed sections from $\mathcal{P}$. Let v_i be the initial vertex of the edge e_i . We introduce new variables $\bar{u}^{(i)} = \{u_{ie} \mid e \notin T, e \notin \mathcal{P}\}$, $\bar{z}^{(i)} = \{z_{ie} \mid e \notin T, e \notin \mathcal{P}\}$ for $1 \leq i \leq m$, as follows

$$u_{ie} = h(r(v_0, v_i))^{-1} h(c_e) h(r(v_0, v_i)), \quad (8)$$

$$h(e_i)^{-1} u_{ie} h(e_i) = z_{ie}. \quad (9)$$

Without loss of generality we can assume $v_0 = (1)$.

Lemma 16. *Let Ω' be periodized with respect to a periodic structure $\langle \mathcal{P}, R \rangle$. Let K be the subgroup of $G_{\Omega'}$ generated by*

$$Y_0 = \{\bar{t}, \{h(e) \mid e \in T_0\}, h(\bar{c}^{(1)}), \bar{u}^{(i)}, \bar{z}^{(i)}, i \in [1, m]\}.$$

If $|\bar{c}^{(2)}| = s \geq 1$ then the group $G_{\Omega'}$ splits as a fundamental group of a graph of groups with two vertices, where one vertex group is K and the other is a free abelian group generated by $h(\bar{c}^{(2)}), h(\bar{c}^{(1)})$. The corresponding edge group is generated by $h(\bar{c}^{(1)})$. The other edges are loops at the vertex with vertex group K , have stable letters $h(e_i)$, $i \in [1, m]$, and associated subgroups $\langle \bar{u}^i \rangle$, $\langle \bar{z}^i \rangle$. If $\bar{c}^{(2)} = \emptyset$ then there is no vertex with abelian vertex group.

Proof. We are going to study in more detail how the variables $h(e_i)$, $i \in [1, m]$ can participate in the equations from Ω' rewritten in the set of variables Y .

If h_k does not lie on a closed section from $\mathcal{P}$, or $h_k \notin \mathcal{P}$, but $e \in T$ (where $h(e) = h_k$), then h_k belongs to the basis Y and is distinct from each of $h(e_1), \dots, h(e_m)$. Now let $h(e) = h_k$, $h_k \notin \mathcal{P}$ and $e \notin T$. Then $e = r_1 c_e r_2$, where r_1, r_2 are paths in T . Since $e \in \Gamma_0$, $h(c_e)$ belongs to $\langle c^{(1)} \rangle$ modulo commutation of cycles. The vertices (k) and $(k+1)$ lie in the same connected component of the graph Γ_0 and, hence, they are connected by a path s in the forest T_0 . Furthermore, r_1 and $s r_2^{-1}$ are paths in the tree T connecting the vertices (k) and v_0 . Consequently, $r_1 = s r_2^{-1}$. Thus, $e = s r_2^{-1} c_e r_2$ and $h_k = h(s) h(r_2)^{-1} h(c_e) h(r_2)$. The variable $h(e_i)$, $i \in [1, m]$ can occur in the right-hand side of the expression obtained (written in the basis Y) only in $h(r_2)$ and at most once. Moreover, the sign of this occurrence (if it exists) depends only on the orientation of the edge e_i with respect to the root v_0 of the tree T . If $r_2 = r'_2 e_i^{\pm 1} r''_2$ then all the occurrences of the variable $h(e_i)$ in the words h_k written in the basis Y , with $h_k \notin \mathcal{P}$, are contained in the occurrences of words of the form $h(e_i)^{\mp 1} h((r'_2)^{-1} c_e r'_2) h(e_i)^{\pm 1}$, that is, in occurrences of the form $h(e_i)^{\mp 1} h(c) h(e_i)^{\pm 1}$, where c is a certain cycle of the graph Γ starting at the initial vertex of the edge $e_i^{\pm 1}$.

Therefore all the occurrences of $h(e_i)$, $i \in [1, m]$ in the equations corresponding to $\lambda \notin \mathcal{P}$ are of the form $h(e_i^{-1}) h(c) h(e_i)$. Also, $h(e_i)$ does not occur in the equations corresponding to $\lambda \in \mathcal{P}$ in the basis Y . The system Ω' is equivalent to the following system in the variables Y :

$$u_{ie} = h(r(v_0, v_i))^{-1} h(c_e) h(r(v_0, v_i)),$$

$$\begin{aligned}
h(e_i)^{-1} u_{ie} h(e_i) &= z_{ie}, \\
[u_{ie_1}, u_{ie_2}] &= 1, \\
[h(c_1), h(c_2)] &= 1, \quad c_1, c_2 \in c^{(1)}, c^{(2)}, \\
\bar{\psi}(Y_0) &= 1,
\end{aligned}$$

where $\bar{\psi}(Y_0)$ does not contain $h(e_i)$, $\bar{c}^{(2)}$. Let $K = G_{\bar{\psi}}$. Then to obtain G_Ω we first take an HNN-extension of the group K with abelian associated subgroups generated by $\bar{u}^{(i)}$ and $\bar{z}^{(i)}$ and stable letters $h(e_i)$, and then extend the centralizer of the image of $\langle \bar{c}^{(1)} \rangle$ by the free abelian subgroup generated by the images of $\bar{c}^{(2)}$. $\square$

Proposition 3 now follows from Lemma 16.

(d) Abelian splitting: long shift. If Ω does not satisfy the conditions of (a)–(c), we perform QH-shortening, then apply the entire transformation and then, if possible, the transformation (D7).

Lemma 17. *Let*

$$v_1 \rightarrow v_2 \rightarrow \cdots \rightarrow v_r \rightarrow \cdots$$

be an infinite path in $T(\Omega)$. Then there exists a natural number N such that all the generalized equations in vertices v_n , $n \geq N$ satisfy the general JSJ-case (d).

Proof. Indeed, the Tietze cleaning either replaces the group by its proper free factor or decreases the complexity. Every time when the case (a) holds we replace G by some vertex group in a non-trivial abelian splitting of G . This can be done only finitely many times [12]. Every time when case (c) takes place, we decrease the complexity. $\square$

Proposition 4. *The general JSJ case (d) cannot be repeated infinitely many times.*

Proof. Consider an infinite path

$$r = v_1 \rightarrow v_2 \rightarrow \cdots \rightarrow v_m \cdots \quad (10)$$

We have $\tau'_{v_i} = \tau'_v$.

Denote by μ_i the carrier base of the equation Ω_{v_i} . The path (10) will be called μ -reducing if $\mu_1 = \mu$, and either μ_2 does not overlap with its double or they overlap but $|\mu| \leq 2|\alpha(\mu) - \alpha(\bar{\mu})|$, and μ occurs in the sequence $\mu_1, \dots, \mu_{m-1}$ at least twice.

The path (10) is called *prohibited* if it can be represented in the form

$$r = r_1 s_1 \cdots r_l s_l r', \quad (11)$$

such that for some sequence of bases $\eta_1, \dots, \eta_l$ the following three properties hold:

- (1) every base occurring at least once in the sequence $\mu_1, \dots, \mu_{m-1}$ occurs at least $4n(1 + f_1(\Omega_{v_2}))$ times in the sequence $\eta_1, \dots, \eta_l$, where n is the number of pairs of bases in equations Ω_{v_i} ,
- (2) the path r_i is η_i -reducing,
- (3) every transfer base of some equation of path r is a transfer base of some equation of path r' .

Every infinite path contains a prohibited subpath. Indeed, let ω be the set of all bases occurring in the sequence $\mu_1, \dots, \mu_m, \dots$ infinitely many times, and $\tilde{\omega}$ the set of all bases, that are transfer bases of infinitely many equations Ω_{v_i} . If one cuts out some finite part in the beginning of this infinite path, one can suppose that all the bases in the sequence $\mu_1, \dots, \mu_m, \dots$ belong to ω and each base that is a transfer base of at least one equation, belongs to $\tilde{\omega}$. Such an infinite path for any $\mu \in \omega$ contains infinitely many non-intersecting μ -reducing finite subpaths. Hence, it is possible to construct a subpath (11) of this path satisfying the first two conditions in the definition of a prohibited subpath. Making r' longer one obtains a prohibited subpath.

Suppose we have a prohibited path (10). As before, let μ_i denote the carrier base of Ω_{v_i} , and $\omega = \{\mu_1, \dots, \mu_{m-1}\}$ and $\tilde{\omega}$ denote the set of such bases which are transfer bases for at least one equation in (10). By ω_1 denote the set of such bases μ for which either μ or $\bar{\mu}$ belongs to $\omega \cup \tilde{\omega}$. By ω_2 denote the set of all the other bases. Let

$$\alpha(\omega) = \min(\min_{\mu \in \omega_2} \alpha(\mu), j),$$

where j is the boundary between active and non-active sections.

Let

$$X_\mu \doteq U[\alpha(\mu), \beta(\mu)].$$

If (Ω_{v_i}, U) corresponds to a vertex in the sequence (10) then denote

$$|U_\omega| = \sum_{i=1}^{\alpha(\omega)-1} |U_i|, \quad (12)$$

$$\psi_\omega(U) = \sum_{\mu \in \omega_1} |X_\mu| - 2|U|_\omega. \quad (13)$$

We call $\psi_\omega(U)$ the *excess* of Ω . Since the case (a) is not applicable, $\psi_\omega(U) \geq 0$ and is comparable with the length of the section $[1, \alpha(\omega)]$. Consider the quadratic part of $\tilde{\Omega}_{v_1}$ which is situated to the left of $\alpha(\omega)$. Let A be the length of the quadratic part and B be the length of the non-quadratic part.

Lemma 18. *Suppose there exists a function $f_1(\Omega_{v_1})$ such that we have an inequality*

$$A \leq B f_1. \quad (14)$$

Then there is a function $f_2(\Omega_{v_1})$ such that

$$|U_\omega| \leq \psi_\omega(U) f_2. \quad (15)$$

Proof. $|U_\omega| = A + B \leq (1 + f_1)B \leq (1 + f_1)\psi_\omega(U)$, because $B \leq \psi_\omega(U)$. $\square$

From the definition of the process it follows that all the words $U^{(i)}[1, \rho_i + 1]$ are the ends of the word $U^{(1)}[1, \rho_1 + 1]$, that is

$$U^{(1)}[1, \rho_1 + 1] \doteq u_i U^{(i)}[1, \rho_i + 1]. \quad (16)$$

On the other hand any base $\mu \in \omega_2$ participates in these transformations neither as a carrier base, nor as a transfer base. Hence, $U^{(1)}[\alpha(\omega), \rho_1 + 1]$ is the end of the word $U^{(i)}[1, \rho_i + 1]$, that is,

$$U^{(i)}[1, \rho_i + 1] \doteq v_i U^{(1)}[\alpha(\omega), \rho_1 + 1]. \quad (17)$$

So we have

$$|U_\omega^{(i)}| - |U_\omega^{(i+1)}| = |v_i| - |v_{i+1}| = |u_{i+1}| - |u_i| = |X_{\mu_i}^{(i)}| - |X_{\mu_i}^{(i+1)}|. \quad (18)$$

In particular (13), (18) imply that $\psi_\omega(U^{(1)}) = \psi_\omega(U^{(2)}) = \dots = \psi_\omega(U^{(m)}) = \psi_\omega$. Denote the number (18) by δ_i .

Let the path (10) be μ -reducing. Estimate $d(U_m) = \sum_{i=1}^{m-1} \delta_i$ from below. First notice that if $\mu_{i_1} = \mu_{i_2} = \mu$ for $i_1 < i_2$ and $\mu_i \neq \mu$ for $i_1 < i < i_2$ then

$$\sum_{i=i_1}^{i_2-1} \delta_i \geq |U^{i_1+1}[1, \alpha(\bar{\mu}_{i_1+1})]|. \quad (19)$$

Indeed, if $i_2 = i_1 + 1$ then $\delta_{i_1} = d(U^{(i_1)}[1, \alpha(\bar{\mu})]) = d(U^{(i_1+1)}[1, \alpha(\bar{\mu})])$. If $i_2 > i_1 + 1$ then $\mu_{i_1+1} \neq \mu$ and μ is a transfer base in the equation $\Omega_{v_{i_1+1}}$. Hence, $\delta_{i_1+1} + |U^{(i_1+2)}[1, \alpha(\mu)]| = |U^{(i_1+1)}[1, \alpha(\mu_{i_1+1})]|$. Now (19) follows from

$$\sum_{i=i_1+2}^{i_2-1} \delta_i \geq |U^{(i_1+2)}[1, \alpha(\mu)]|.$$

So, if the bases μ_2 and $\bar{\mu}_2$ do not intersect in the equation Ω_{v_2} then (19) implies that

$$\sum_{i=1}^{m-1} \delta_i \geq |U^{(2)}[1, \alpha(\bar{\mu}_2)]| \geq |X_{\mu_2}^{(2)}| \geq |X_\mu^{(2)}| = |X_\mu^{(1)}| - \delta_1,$$

which implies that

$$\sum_{i=1}^{m-1} \delta_i \geq \frac{1}{2} |X_\mu^{(1)}|. \quad (20)$$

We obtain the same inequality if μ_2 overlaps with $\bar{\mu}_2$ but $|\mu_2| \leq 2|\alpha(\mu_2) - \alpha(\bar{\mu}_2)|$.

Suppose now that the path (10) is prohibited. Hence, it can be represented in the form (11). From the definition (13) we have $\sum_{\mu \in \omega_1} d(X_\mu^{(m)}) \geq \psi_\omega$. So, at least for one base $\mu \in \omega_1$ the inequality $d(X_\mu^{(m)}) \geq \frac{1}{2n} \psi_\omega$ holds. Because $X_\mu^{(m)} \doteq (X_{\bar{\mu}}^{(m)})^{\pm 1}$, we can suppose that $\mu \in \omega \cup \tilde{\omega}$. Let m_1 be the length of the path $r_1 s_1 \dots r_l s_l$ in (11). If $\mu \in \tilde{\omega}$ then by the third part of the definition of a prohibited path there exists $m_1 \leq i \leq m$ such that μ is a transfer base of Ω_{v_i} . Hence, $d(X_{\mu_i}^{(m_1)}) \geq d(X_{\mu_i}^{(i)}) \geq d(X_\mu^{(i)}) \geq d(X_\mu^{(m)}) \geq \frac{1}{2n} \psi_\omega$. If $\mu \in \omega$ then take μ instead of μ_i . We proved the existence of a base $\mu \in \omega$ such that

$$|X_\mu^{(m_1)}| \geq \frac{1}{2n} \psi_\omega. \quad (21)$$

By the definition of a prohibited path, the inequality (20) and the inequality $d(X_\mu^{(i)}) \geq d(X_\mu^{(m_1)})$ ($1 \leq i \leq m_1$), we obtain

$$\sum_{i=1}^{m_1-1} \delta_i \geq \max \left\{ \frac{1}{4n} \psi_\omega, 1 \right\} (4n f_2). \quad (22)$$

By (18) the sum in the left part of the inequality (22) equals $d_\omega(U^{(1)}) - d_\omega(U^{(m_1)})$. Hence,

$$|U_\omega^{(1)}| \geq \max \left\{ \frac{1}{4n} \psi_\omega, 1 \right\} (4n f_2) = \psi_\omega f_2,$$

which, in case of the inequality (14), contradicts (15). This contradiction was obtained from the supposition that there are prohibited paths (10). Hence there are no prohibited paths. The proposition is proved. $\square$

Proof of Theorem 1. We perform the elimination process for the generalized equation $\Omega = \Omega_{v_0}$ corresponding to G . If the process goes infinitely we obtain one of the following:

- free splitting of G with at least on non-trivial free factor (4.4.1) and, maybe, some surface group factor (4.4.2),
- a decomposition of G as the fundamental group of a graph of groups with QH-vertex groups corresponding to quadratic sections (4.4.4 (a)),
- decomposition of G as the fundamental group of a graph of groups with abelian vertex groups corresponding to periodic structures (see Proposition 3 in 4.4.4 (c)) and HNN-extensions with stable letters infinitely longer than the generators of the abelian associated subgroups (4.4.4 (c)).

Then we continue the elimination process with the generalized equation Ω_{v_0} where the active part corresponds to weakly rigid subgroups. In the case of an HNN-extension the complexity $\tau(\Omega_{v_1})$ is smaller than $\tau(\Omega_{v_0})$. In the other cases the Delzant–Potyagailo complexity of weakly rigid subgroups is smaller by Proposition 2. Therefore this procedure stops. At the end we obtain some generalized equation $\Omega_{v_{fin}}$ with all non-active sections. Continuing the elimination process till the end, we obtain a complete system of linear equations with integer coefficients $\Sigma_{complete}$ on the lengths of items h_i 's that is automatically satisfied, therefore the associated maximal abelian subgroups are length-isomorphic.

Notice that if $G_{i+1} = G_{\Omega_{v_{i+1}}}$ is obtained as the fundamental group of a graph of groups with two vertices, where G_i is one vertex group and the other vertex is a QH-vertex, then one can also obtain G_{i+1} from G_i by taking a free product of G_i and a free group and then the HNN-extension with cyclic associated subgroups. If G_{i+1} is an amalgamated product of G_i and a free abelian group then it can be obtained from G_i as a series of extensions of centralizers.

This completes the proof of Theorem 1. □

Remark 5. *If we begin with the group $\tilde{G}$ mentioned in Remark 2 with free but not necessary regular length function in Λ then in the Elimination process we work with the generalized equation Ω and add a finite number of elements from $\tilde{G}$. Thus we got an embedding of $\tilde{G}$ in a group that can be represented as a union of a finite series of groups*

$$G_1 < G_2 < \cdots < G_n = G,$$

where

1. G_1 is a free group,
2. G_{i+1} is obtained from G_i by finitely many HNN-extensions in which associated subgroups are maximal abelian and length isomorphic as subgroups of Λ .

Remark 6. *As a result of the Elimination process, the equation $\Omega_{v_{fin}}$ (we will denote it Ω_{fin}) is defined on the multi-interval I , that is, a union of closed sections which have a natural hierarchy: a section σ_1 is smaller than a section σ_2 if the largest base on σ_2 is infinitely larger than the largest base on σ_1 .*

The lengths of bases satisfy the system of linear equations $\Sigma_{complete}$.

5 Proof of Theorems 3 and 4

Suppose G , as above, is a Λ -free group with a regular action. Let Ω be a generalized equation for G corresponding to the union of closed sections I , $G = \langle \mathcal{M} \mid \Omega(\mathcal{M}) \rangle$. Consider the Cayley graph $X = \text{Cay}(G, \mathcal{M})$ of G with respect to the generators $\mathcal{M}$. Assign to edges of $\text{Cay}(G, \mathcal{M})$ their lengths in Λ and consider edges as closed intervals in Λ of the corresponding length. For each relation

between bases of Ω , $\lambda_{i_1} \cdots \lambda_{i_k} = \lambda_{j_1} \cdots \lambda_{j_m}$ (without cancelation) there is a loop in X labeled by this relation. Then the path on the boundary of this loop labeled by $\lambda_{i_1} \cdots \lambda_{i_k}$ has the same length in Λ as the path labeled by $\lambda_{j_1} \cdots \lambda_{j_m}$. If x is a point on the path $\lambda_{i_1} \cdots \lambda_{i_k}$ at the distance $d \in \Lambda$ from the beginning of the path, and x' is a point on the path $\lambda_{j_1} \cdots \lambda_{j_m}$ at the distance d from the beginning, then we say that x and x' are in the same leaf. In other words, after we substitute generators in $\mathcal{M}$ by their infinite word representations, we “fold” loops into segments. We consider the equivalence relation between points of X generated by all such pairs $x \sim x'$. Equivalence classes of this relation are called *leaves*. We also glue an arc isometric to a unit interval between each x and x' . Let $\mathcal{F}$ be the foliation (the set of leaves). One can define a foliated complex $\Sigma = \Sigma(X, \mathcal{F})$ associated to X as a pair $(X, \mathcal{F})$. We define a metric space $T(X)$ as follows. The paths in Σ can travel vertically (along the leaves) and horizontally (along the intervals in Λ). The length of a path γ in $\Sigma(X, \mathcal{F})$ (denoted $\|\gamma\|$) is the sum of the lengths of horizontal intervals in γ . The pseudo-distance d between two points of Σ is defined as the length of a minimal path between them. A metric space T is obtained from Σ by identifying points at pseudo-distance 0. We extend naturally the left action of G on X to the action on T .

Lemma 19. (1) T is a Λ -tree.

(2) The left action of G on T is free.

Proof. We will prove the second statement first. Suppose x is a vertex of X , $g \in G$. Then gx is also a vertex of X . If $d(gx, x) = 0$ in Σ , then x and gx are in the same leaf. Hence, there is a vertical path $x \sim x_1 \sim \cdots \sim x_k = gx$ that uses relations from $\Omega(\mathcal{M})$

$$R_1 : p_{11} q_{11} = p_{12} \mu_1 q_{12}, \quad R_2 : p_{21} \mu_1 q_{21} = p_{22} \mu_2 q_{22}, \quad \dots, \quad R_k : p_{k1} \mu_{k-1} q_{k1} = p_{k2} q_{k2},$$

where all p_{ij}, q_{ij} are products of some bases (G is generated by bases from $\mathcal{M}$); x is a vertex between p_{11} and q_{11} , x_i is a vertex on μ_i , x_k is a vertex between p_{k2} and q_{k2} . Then $\mu_i = \mu_{i1}\mu_{i2}$, where

$$\ell(p_{11}) = \ell(p_{12} \mu_{11}), \quad \ell(p_{21} \mu_{11}) = \ell(p_{22} \mu_{21}), \dots, \ell(p_{i1} \mu_{(i-1)1}) = \ell(p_{i2} \mu_{i1}), \quad \ell(p_{k1} \mu_{(k-1)1}) = \ell(p_{k2}).$$

Since relations are coming from the generalized equation Ω , p_{11} and $p_{12}\mu_{11}$ are the same infinite words, as well as $p_{i1} \mu_{(i-1)1}$ and $p_{i2} \mu_{i1}$ are the same infinite words for $i = 2, \dots, k-1$, and $p_{k1} \mu_{(k-1)1}$ and p_{k2} are the same infinite words. Therefore $\mu_{11} = p_{12}^{-1} p_{11} = p_{21}^{-1} p_{22} \mu_{21}$, $\mu_{21} = p_{31}^{-1} p_{32} \mu_{31}, \dots$, $\mu_{(k-1)1} = p_{k1}^{-1} p_{k2}$ and $p_{12}^{-1} p_{11} = p_{21}^{-1} p_{22} \dots p_{k1}^{-1} p_{k2}$.


Then $p_{k2}^{-1} p_{k1} \cdots p_{12}^{-1} p_{11} = 1$ in G . But the element $p_{k2}^{-1} p_{k1} \cdots p_{12}^{-1} p_{11} \in G$ takes $gx \in X$ to $x \in X$, hence $p_{k2}^{-1} p_{k1} \cdots p_{12}^{-1} p_{11}gx = x$. Therefore $gx = x$ and $g = 1$.

If x is not a vertex of X , then x is a point on some edge of X labeled by μ_0 and joining some x_0 and $x_0\mu_0$, while gx , by translation in the Cayley graph, is a point on the edge of X labeled by μ_0 and joining gx_0 and $gx_0\mu_0$. There is a vertical path $x \sim x_1 \sim \cdots \sim x_k = gx$ that uses relations of $\Omega(\mathcal{M})$

$$R_1 : p_{11} \mu_0 q_{11} = p_{12} \mu_1 q_{12}, \quad R_2 : p_{21} \mu_1 q_{21} = p_{22} \mu_2 q_{22}, \quad \dots, \quad R_k : p_{k1} \mu_{k-1} q_{k1} = p_{k2} \mu_0 q_{k2}.$$

Let y be a vertex of X belonging to one of the loops $R_1, \dots, R_k$ and nearest to the leaf joining x and gx . Suppose y belongs to the loop R_2 and it is a vertex between p_{21} and μ_1 . Then there is a leaf joining y and gy that uses the relations $R_2, \dots, R_k, R_1$, and we can show that $g = 1$ as above.

Now we prove that T is a Λ -tree. Suppose, we have a loop γ in Σ beginning at x , which is a vertex of X ($x \in VX$). We are going to show that the image of this loop becomes degenerate in T (if we remove all subpaths of the form pp^{-1} from the image of γ we get a point). Notice that the image of any closed horizontal path in Σ is degenerate in T , so in the loop we can replace a horizontal subpath joining any two vertices $y, z \in VX$ by any other horizontal path joining y and z . Represent $\gamma = \gamma_1\gamma_2$, where γ_1 is a sequence of vertical and horizontal paths between x and another vertex y

Figure 8: Replacing γ_1 by γ_3

of X , and γ_2 is a horizontal path between y and x such that if there is a horizontal subpath p at the end of γ_1 then p does not contain another vertex $z \in VX$. We can use induction on the number of vertical components in γ_1 , the case of zero vertical components being trivial. We can replace the path γ_1 by a path γ_3 of equal or shorter length with the same image in T such that each vertical subpath of γ_3 is passing through some vertex of X (see Figure 8).




Figure 9: Getting rid of a vertical subpath

Let $z_1 \in VX$ be the first vertex of γ_2 . Suppose the last vertical part of γ_3 begins with z_5 , ends with z_4 and is passing through $z_2 \in VX$. Then the image in T of the circle at z_2 consisting of the vertical path from z_2 to z_4 (see Figure 9) and the corresponding horizontal path from z_4 to z_2 (denote it γ_4) is degenerate. The image of the circle at z_2 consisting of the vertical path from z_2 to z_5 and the horizontal path z_2, z_3, z_5 is also degenerate. Denote by γ_5 the horizontal path between z_3

and z_2 . We now replace γ_2 by $\gamma_5\gamma_4^{-1}\gamma_2$ and γ_1 by the part of γ_1 from x to z_3 . The image of this path in T is degenerate by induction because it has less vertical subpaths. Therefore, T is a Λ -tree. $\square$

Similarly, we can construct a Λ -tree where G acts freely beginning not with the original generalized equation Ω but with a generalized equation obtained from Ω by the application of the Elimination process. In this case we may have closed sections with some items h_i with $\gamma(h_i) = 1$. Then to construct the Cayley graph of G in the new generators we cover items h_i with $\gamma(h_i) = 1$ by bases without doubles. (These bases were previously removed at some stages of the Elimination process as matching pairs).

Proof of Theorem 3. We begin by considering the union of the smallest closed sections, see Remark 6. Denote the union of these sections by σ . The group H of the generalized equation corresponding to their union is a free product of free groups, free abelian groups and closed surface groups. Notice that for those sections for which $\gamma(h_i) = 1$ for some maximal height items, these items h_i are also products of some bases because initially every item is a product of bases. We can assume the following:

1. For all closed sections of σ such that $\gamma(h_i) = 2$ for all items of the maximal height, the number of bases of maximal height cannot be decreased using entire transformation or similar transformation applied from the right of the section (*right entire transformation*);
2. For all closed sections of σ where we apply linear elimination, the number of bases of maximal height cannot be decreased using the transformations as above as well as transformation (E2) (transfer) preceded by creation of necessary boundary connections as in (E5) (denote it (E2₅)), (E3) and first two types of linear elimination (D6) (denote it (D6_{1,2}));

Indeed, if we can decrease the number of bases of maximal height using the transformations described above then we just do this and continue. Since these transformations do not change the total number of bases we can also assume that they do not decrease the number of bases of second maximal height etc. We now re-define the lengths of bases belonging to σ in $\mathbb{R}^k$. We are going to show that all components of the length of every base can be made zeros except for the components which appear to be maximal in the lengths of bases from σ .

Let Ω_σ be the generalized equation corresponding to the sections from σ . Let H be the group of the equation $H = \langle \mathcal{M}_\sigma \mid \Omega_\sigma(M) \rangle$.

Denote by Λ_1 the minimal convex subgroup of Λ containing lengths of all bases in σ , and by Λ' a maximal convex subgroup of Λ_1 not containing lengths of maximal height bases in σ (it exists by Zorn's lemma). Then the quotient Λ_1/Λ' is a subgroup of $\mathbb{R}$. Denote by $\hat{\ell}$ the length function in $\mathbb{R}$ on this quotient induced from ℓ . We consider elements of Λ' as infinitesimals. Denote by $\hat{T}$ the $\mathbb{R}$ -tree constructed from T by identifying points at zero distance (see [7], Theorem 2.4.7). Then H acts on $\hat{T}$. Denote by $\|\gamma\|_{\mathbb{R}}$ the induced length of the path γ in $\hat{T}$, and by $d_{\mathbb{R}}(\bar{x}, \bar{y})$ the induced distance.

However, the action of H on $\hat{T}$ is not free. The action is minimal, that is, there is no non-empty proper invariant subtree. Notice that the canonical projection $f : T \rightarrow \hat{T}$ preserves alignment, and the pre-image of the convex set is convex. The pre-image of a point in $\hat{T}$ is an infinitesimal subtree of T .

Lemma 20. *The action of H on $\hat{T}$ is superstable: for every non-degenerate arc $J \subset \hat{T}$ with non-trivial fixator, and for every non degenerate subarc $S \subset J$, one has $\text{Stab}(S) = \text{Stab}(J)$.*

The proof is the same as the proof of Fact 5.1 in [18].

Lemma 21. *Let σ_1 be a closed section on the lowest level corresponding to a closed surface group and the lengths of the bases satisfy some system of linear equations Σ . Then one can define the lengths of bases in σ_1 in $\mathbb{R}^k$.*

Proof. Denote by $\mathcal{M}_{\sigma+}$ the set of bases (on all the steps of the process of entire transformation applied to the lowest level) with non-zero oldest component and by $\mathcal{M}_{\sigma 0}$ the rest of the bases (infinitesimals). Denote by $\hat{\ell}$ the projection of the length function ℓ to Λ_1/Λ' as before. Then $\lambda \in \mathcal{M}_{\sigma+}$ if and only if $\hat{\ell}(\lambda) > 0$. We apply the entire transformation to σ_1 . If we obtain an overlapping pair or an infinitesimal section, where the process goes infinitely, we declare it non-active and move to the right. This either decreases, or does not change the number of bases in the active part. Therefore, we can assume that the process goes infinitely and the number of bases in $\mathcal{M}_{\sigma+}$ never decreases. Therefore bases from $\mathcal{M}_{\sigma 0}$ are only used as transfer bases.

We will show that the stabilizer of a pre-image of a point, an infinitesimal subtree T_0 of T is generated by some elements in $\mathcal{M}_{\sigma 0}$. An element h from H belongs to such a stabilizer if $\hat{\ell}(h) = 0$. If some product of bases not only from $\mathcal{M}_{\sigma 0}$ has infinitesimal length (denote this product by g), then the identity and g belong to leaves at the infinitesimal distance δ in Σ . Therefore using elementary operations we can obtain a base of length δ . Denote by Λ'' the minimal convex subgroup of Λ containing all elements of the same height as bases from $\mathcal{M}_{\sigma 0}$.

This implies the following lemma which we need to finish the proof of lemma 21.

Lemma 22. *Let $\bar{\sigma}_1$ be the projection of the quadratic section σ_1 to $\Lambda_1/(\Lambda'' \cap \Lambda_1)$. Suppose the process of entire transformation for σ_1 goes infinitely and the number of bases in $\mathcal{M}_{\sigma+}$ never decreases. Then the process of entire transformation for $\bar{\sigma}_1$ goes infinitely too and the number of bases in $\mathcal{M}_{\sigma+}$ never decreases.*

If $\ell(g) \in \Lambda'$ then g is a product of bases in $\mathcal{M}_{\sigma 0}$ and $\ell(g) \in \Lambda''$.

This lemma implies that there is no element in H infinitely larger than all bases in $\mathcal{M}_{\sigma 0}$, but infinitely smaller than all bases in $\mathcal{M}_{\sigma+}$.

Therefore, we can make all components of the lengths of bases in $\mathcal{M}$ zeros except for those which are maximal components of some bases in $\mathcal{M}$. Since $\mathcal{M}$ is a finite set, the number of such components is finite, and the length is defined in $\mathbb{R}^k$ for some k not larger than the number of pairs of bases. $\square$

Let σ be a closed section corresponding to infinite linear elimination process for a free group as in Lemma 7 (c). We cover (for technical reasons) items h_i with $\gamma(h_i) = 1$ by bases without doubles. (These bases were removed at some stages of the Elimination process as matching pairs).

Lemma 23. *Let σ be a closed section corresponding to infinite linear elimination process for a free group as in Lemma 7 (c). Let $\bar{\sigma}$ be the projection of section σ to Λ_1/Λ'' . Suppose the process of linear elimination on σ goes infinitely and some generalized equation Ω' (with the minimal possible number of bases) appears in this process infinitely many times. Then the process of linear elimination on $\bar{\sigma}$ goes infinitely. Moreover, suppose for any base μ with $\hat{\ell}(\mu) = 0$ obtained by elementary transformations, $\ell(\mu) \in \Lambda''$. If $H = \langle \mathcal{M}_\sigma \mid \Omega_\sigma(\mathcal{M}_\sigma) \rangle$, and $g \in H, \hat{\ell}(g) = 0$ then $\ell(g) \in \Lambda''$.*

Proof. The first statement holds because the steps that we make in the process depend only on the situation of bases of maximal height. If some product g of bases not only from $\mathcal{M}_{\sigma 0}$ has infinitesimal length, then the identity and g belong to leaves at the infinitesimal distance δ in Σ . Therefore using elementary operations we can obtain a base μ of length comparable to δ . Therefore $\delta \in \Lambda''$. $\square$

Lemma 24. *Let $\bar{\sigma}$ be a closed section corresponding to infinite linear elimination process for a free group. Suppose the process of linear elimination on $\bar{\sigma}$ goes infinitely and some generalized equation Ω (with the minimal possible number of bases) appears in this process infinitely many times. Suppose all items have length in $\mathbb{R}$. Then applying elementary transformations we cannot decrease the difference k between the number of items and the total number of basic and boundary equations. This difference is the rank of the free group that is the coordinate group of Ω .*

Proof. Every equation Ω' that is obtained from Ω by elementary operations has the free group of rank k as the coordinate group. If we cut bases of Ω' along boundary connections and start removing eliminable bases, we have to remove them all. Every time when we remove eliminable base we express one item in terms of the others. There must be k items left at the end. $\square$

Lemma 24 implies that if σ is a closed section corresponding to infinite linear elimination process for a free group, then a base μ with $\hat{\ell}(\mu) = 0$ can only be obtained by elementary transformations from bases in $\mathcal{M}_{\sigma 0}$. Therefore, by Lemma 23, if $H = \langle \mathcal{M}_{\sigma} \mid \Omega_{\sigma}(\mathcal{M}_{\sigma}) \rangle$, and $g \in H$, $\hat{\ell}(g) = 0$ then $\ell(g) \in \Lambda''$.

Finally, if a finitely generated abelian group H acts freely on a Λ -tree, then it is isometrically embedded into Λ (see, for example, [3]).

Lemma 25. ([7], Lemma 1.1.6) *If H is a finitely generated ordered abelian group, it has a decomposition $H = \Lambda_1 \oplus \dots \oplus \Lambda_k$ for some $k \geq 0$, where the ordering is the lexicographic ordering and $\Lambda_1, \dots, \Lambda_k$ are subgroups of $\mathbb{R}$.*

Therefore, H has a free basis $h_1, \dots, h_k$ such that $ht(h_1) > \dots > ht(h_k)$. We can now make all the components in the length of the basis elements zeros except those corresponding to $ht(h_1), \dots, ht(h_k)$. This defines a free length function in $\mathbb{R}^n$ on H . The new lengths of elements in H satisfy the same system of linear equations as their lengths in Λ .

Using induction on the number of levels obtained in the Elimination process, we similarly prove the statement of Theorem 3.

The points of an $\mathbb{R}^n$ -tree, where G acts freely are the leaves in the foliation corresponding to the new length of bases in $\mathbb{R}^n$. The new lengths of bases are exactly their Lyndon lengths. $\square$

Proof of Theorem 4. Notice that in the case when $\tilde{G}$ is a finitely presented group with a free length function in Λ (not necessary regular) it can be embedded in the group with a free regular length function in Λ by Remark 2. That group can be embedded in $R(\Lambda', X)$. When we make a generalized equation for $\tilde{G}$, we have to add only a finite number of elements from $R(\Lambda', X)$. We run the elimination process for this generalized equation as we did in the proof of Theorem 1 and obtain a group G as in Remark 5, where $\tilde{G}$ is embedded, and then redefine the length of elements of G in $\mathbb{R}^n$ as above. Therefore G acts freely and regularly on a $\mathbb{R}^n$ -tree. The theorem is proved.

Moreover, using Lemmas 22 and 23, and induction one shows that the length function in $\mathbb{R}^n$ and in Λ defined on G is regular. Therefore we proved Theorem 4.

Remark 7. *If $\tilde{G}$ is a finitely presented group which has a free (not necessary regular) length function in Λ , then it acts freely on a Λ -tree, and one can construct, as in [4], a foliated band complex with measured foliation with horizontal measure in Λ .*

References

- [1] R. Alperin and H. Bass, *Length functions of group actions on Λ -trees*. Combinatorial group theory and topology, (Ed. S. M. Gersten and J. R. Stallings), Annals of Math. Studies **111**, 265–378. Princeton University Press, 1987.
- [2] R. Alperin and K. Moss, *Complete trees for groups with a real length function*. J. London Math. Soc. (2) **31**, 1985, 55–68.
- [3] H. Bass *Groups acting on non-archimedean trees*. Arboreal group theory, 1991, 69–130.
- [4] M. Bestvina and M. Feighn, *Stable actions of groups on real trees*. Invent. Math., **121** no. 2 (1995), 287–321.
- [5] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*. Springer, 1999.

- [6] I. Chiswell, *Abstract length functions in groups*. Math. Proc. Cambridge Philos. Soc., **80** no. 3 (1976), 451–463.
- [7] I. Chiswell, *Introduction to Λ -trees*. World Scientific, 2001.
- [8] I. Chiswell, *A-free groups and tree-free groups*. *Algorithms, Languages, Logic* (Ed. A. Borovik), Contemp. Math., Amer. Math. Soc. **378** (2005), 79–86.
- [9] I. Chiswell and T. Muller, *Embedding theorems for tree-free groups*. Under consideration for publication in Math. Proc. Camb. Phil. Soc.
- [10] F. Dahmani, *Existential questions in (relatively) hyperbolic groups*. Israel J. Math **173** (2009), 91–124.
- [11] F. Dahmani and D. Groves, *The Isomorphism Problem for Total Relatively Hyperbolic Groups*. Publ. Math., Inst. Hautes Etudes Sci. **107** no. 1 (2008), 211–290.
- [12] T. Delzant and L. Potyagailo, *Accessibilité hiérarchique des groupes de présentation finie*. (French) Topology **40** no. 3 (2001), 617–629.
- [13] D. Gaboriau, G. Levitt and F. Paulin, *Pseudogroups of isometries of $\mathbb{R}$ and Rips’ Theorem on free actions on $\mathbb{R}$ -trees*. Israel. J. Math., **87** (1994), 403–428.
- [14] D. Gaboriau and G. Levitt and F. Paulin, *Pseudogroups of isometries of $\mathbb{R}$: reconstruction of free actions on $\mathbb{R}$ -trees*. Ergodic Theory Dynam. Systems **15** (1995), 633–652.
- [15] D. Gildenhuys, O. Kharlampovich and A. Myasnikov, *CSA-groups and separated free constructions*. Bull. Austral. Math. Soc., **52** no. 1 (1995), 63–84.
- [16] A. M. W. Glass, *Partially ordered groups*. Series in Algebra, **7**, World Scientific, 1999.
- [17] D. Groves, *Limit groups for relatively hyperbolic groups, II: Makanin-Razborov diagrams*. Geom. Topol. **9** (2005), 2319–2358.
- [18] V. Guirardel, *Limit groups and groups acting freely on $\mathbb{R}^n$ -trees*. Geom. Topol., **8** (2004), 1427–1470.
- [19] N. Harrison, *Real length functions in groups*. Trans. Amer. Math. Soc. **174** (1972), 77–106.
- [20] A. H. M. Hoare, *On length functions and Nielsen methods in free groups*. J. London Math. Soc. (2) **14** (1976), 188–192.
- [21] A. H. M. Hoare, *Nielsen method in groups with a length function*. Math. Scand. **48** (1981), 153–164.
- [22] B. Khan, A. Myasnikov and D. Serbin, *On positive theories of groups with regular free length functions*. Internat. J. Algebra and Comput. **17** no. 1 (2007), 1–26.
- [23] O. Kharlampovich and A. Myasnikov, *Irreducible affine varieties over a free group. II: Systems in triangular quasi-quadratic form and description of residually free groups*. J. of Algebra **200** no. 2 (1998), 517–570.
- [24] O. Kharlampovich and A. Myasnikov, *Implicit function theorems over free groups*. J. of Algebra **290** no. 1 (2005), 1–203.
- [25] O. Kharlampovich, A. Myasnikov, V. Remeslennikov and D. Serbin *Groups with free regular length functions in $\mathbb{Z}^n$* . ArXiv:0907.2356, Trans. Amer. Math. Soc., in print.

- [26] O. Kharlampovich, A. Myasnikov and D. Serbin, *Regular completions of $\mathbb{Z}^n$ -free groups*. Preprint, 2011.
- [27] V. Kopytov and N. Medvedev, *Right-ordered groups*. Siberian School of Algebra and Logic. Consultants Bureau, New York, 1996.
- [28] R. Lyndon, *Length functions in groups*. Math. Scand. **12** (1963), 209–234.
- [29] R. C. Lyndon and P. E. Schupp, *Combinatorial group theory*. Ergebnisse der Mathematik und ihrer Grenzgebiete **89**, Springer-Verlag, Berlin, Heidelberg, New York, 1977.
- [30] G. S. Makanin, *Equations in a free group*. (Russian), Izv. Akad. Nauk SSSR, Ser. Mat., **46** (1982), 1199–1273 (transl. in Math. USSR Izv. **21** (1983)).
- [31] A. Myasnikov and V. Remeslennikov, *Length functions on free exponential groups*. Proc. N 26. IITPM SO RAN, Omsk, 1996, 1–34.
- [32] A. Myasnikov, V. Remeslennikov and D. Serbin, *Regular free length functions on Lyndon’s free $\mathbb{Z}[t]$ -group $F^{\mathbb{Z}[t]}$* , *Algorithms, Languages, Logic* (Ed. A. Borovik), Contemp. Math., Amer. Math. Soc. **378** (2005), 37–77.
- [33] J. Morgan and P. Shalen, *Valuations, Trees, and Degenerations of Hyperbolic Structures, I*. Annals of Math, 2nd Ser., **120** no. 3. (1984), 401–476.
- [34] D. Promislow, *Equivalence classes of length functions on groups*. Proc. London Math. Soc (3) **51** (1985), 449–477.
- [35] A. Razborov, *On systems of equations in a free group*. Math. USSR Izvestiya **25** no. 1 (1985), 115–162.
- [36] J.-P. Serre, *Trees*. New York, Springer, 1980.
- [37] A. Schrijver, *Theory of linear and integer programming*. New-York, John Wiley and Sons, 1998.
- [38] D. Y. Rebbeci, *Algorithmic properties of relatively hyperbolic groups*, PhD Thesis, Univ. California, Davis, 2001. <http://front.math.ucdavis.edu/math.GR/0302245>.
- [39] D. Wise The structure of groups with a quasiconvex hierarchy, preprint